

NATIONALE PRAKTIJKRICHTLIJN VOOR DE AANPAK VAN GELOKALISEERDE PROSTAATKANKER: DEEL 1





Het Federaal Kenniscentrum voor de Gezondheidszorg

Het Federaal Kenniscentrum voor de Gezondheidszorg is een parastatale, opgericht door de programmawet (1) van 24 december 2002 (artikelen 259 tot 281) die onder de bevoegdheid valt van de Minister van Volksgezondheid en Sociale Zaken. Het Centrum is belast met het realiseren van beleidsondersteunende studies binnen de sector van de gezondheidszorg en de ziekteverzekering.

Raad van Bestuur

	Effectieve Leden	Plaatsvervangende Leden
Voorzitter	Pierre Gillet	
Leidend ambtenaar RIZIV (vice-voorzitter)	Jo De Cock	Benoît Collin
Voorzitter FOD Volksgezondheid (vice-voorzitter)	Dirk Cuypers	Chris Decoster
Voorzitter FOD Sociale Zekerheid (vice-voorzitter)	Frank Van Massenhove	Jan Bertels
Administrateur-generaal FAGG	Xavier De Cuyper	Greet Musch
Vertegenwoordigers Minister van Volksgezondheid	Bernard Lange	François Perl
	Marco Schetgen	Annick Poncé
Vertegenwoordigers Minister van Sociale Zaken	Olivier de Stexhe	Karel Vermeyen
	Ri De Ridder	Lambert Stamatakis
Vertegenwoordigers Ministerraad	Jean-Noël Godin	Frédéric Lernoux
	Daniel Devos	Bart Ooghe
Intermutualistisch Agentschap	Michiel Callens	Frank De Smet
	Patrick Verertbruggen	Yolande Husden
	Xavier Brenez	Geert Messiaen
Beroepsverenigingen van de artsen	Marc Moens	Roland Lemyé
	Jean-Pierre Baeyens	Rita Cuypers
Beroepsverenigingen van de verpleegkundigen	Michel Foulon	Ludo Meyers
	Myriam Hubinon	Olivier Thonon
Ziekenhuisfederaties	Johan Pauwels	Katrien Kesteloot
	Jean-Claude Praet	Pierre Smiets
Sociale partners	Rita Thys	Leo Neels
	Paul Palsterman	Celien Van Moerkerke
Kamer van Volksvertegenwoordigers	Lieve Wierinck	



Controle

Regeringscommissaris

Yves Roger

Directie

Algemeen Directeur

Raf Mertens

Programmadirectie

Christian Léonard
Kristel De Gauquier

Contact

Federaal Kenniscentrum voor de Gezondheidszorg (KCE)

Doorbuilding (10^e verdieping)

Kruidtuinlaan 55

B-1000 Brussel

Belgium

T +32 [0]2 287 33 88

F +32 [0]2 287 33 85

info@kce.fgov.be

<http://www.kce.fgov.be>

NATIONALE PRAKTIJKRICHTLIJN VOOR DE AANPAK VAN GELOKALISEERDE PROSTAATKANKER: DEEL 1

FRANÇOISE MAMBOURG, PASCALE JONCKHEER, JULIEN PIÉRART, HANS VAN BRABANDT



COLOFON

Titel:	Nationale Praktijkrichtlijn voor de aanpak van gelokaliseerde prostaatkanker: deel 1
Auteurs:	Françoise Mambourg, Pascale Jonckheer, Julien Piérart, Hans Van Brabandt
Reviewers:	Leen Verleye, Jo Robays
Guideline Development Group:	Gert De Meerleer (Radiotherapeut, UZ Gent), Louis Denis (Directeur van « Wij Ook/US TOO »), Axel Feyaerts (Uroloog, Cliniques Universitaires Saint Luc), Laurette Renard (Radiotherapeut, Cliniques Universitaires Saint Luc), Sandrine Rorive (Pathologist, Hôpital Erasme), Dirk Schrijvers (Oncoloog, ZNA Middelheim), Alain Servaes (Patiëntvertegenwoordiger), Bram Spinnewijn (Huisarts, Domus Medica), Bertrand Tombal (Uroloog, Cliniques Universitaires Saint Luc), Roland Van Velthoven (Uroloog, Institut J.Bordet), Geert Villeirs (Radioloog, UGent).
Externe Validatoren:	Nicolas Mottet (Uroloog, CHU-St. Etienne-France), Michel Roland (Huisarts, CEBAM), Guy Soete (Radiotherapeut, UZ Brussel).
Stakeholders:	Thomas Adams (Uroloog, Sint-Augustinus Ziekenhuis), Karin Haustermans (Radiotherapeut, UZ Leuven), Steven Joniau (Uroloog, UZ Leuven), Sara Junius (Radiotherapeut, KU Leuven), Louis Libbrecht (Pathologist, UGent), Raymond Oyen (Radioloog, KU Leuven), Sylvie Rottey (Oncoloog, UZ Gent), Thierry Roumeguere (Uroloog, Hôpital Erasme), Denis Schallier (Radiotherapeut, UZ Brussel)
Belangenconflict:	<p>Honoraria of een andere compensatie voor het schrijven van een publicatie of het deelnemen aan de ontwikkeling ervan: Bram Spinnewijn (Domus Medica aanbeveling)</p> <p>Een beurs, honoraria of fondsen voor een personeelslid of een andere vorm van compensatie voor het uitvoeren van onderzoek: Thomas Adams</p> <p>Consultancy of andere betaalde dienstverlening voor een organisatie die financieel kan winnen of verliezen door de resultaten van dit rapport: Alain Servaes (werkt in studiedienst Euromut; deelname is privé, als vertegenwoordiger patiëntenorganisatie)</p> <p>Betalingen om te spreken, opleidingsvergoedingen, reisondersteuning of betaling voor deelname aan een symposium: Thomas Adams, Bram Spinnewijn (lezing voor huisartsen), Geert Villeirs</p>
Omslagfoto's:	Rechterfoto: Joe Moore
Layout:	Ine Verhulst, Sophie Vaes

**Disclaimer:**

- De externe experts werden geraadpleegd over een (preliminaire) versie van het wetenschappelijke rapport. Hun opmerkingen werden tijdens vergaderingen besproken. Zij zijn geen coauteur van het wetenschappelijke rapport en gingen niet noodzakelijk akkoord met de inhoud ervan.
- Vervolgens werd een (finale) versie aan de validatoren voorgelegd. De validatie van het rapport volgt uit een consensus of een meerderheidsstem tussen de validatoren. Zij zijn geen coauteur van het wetenschappelijke rapport en gingen niet noodzakelijk alle drie akkoord met de inhoud ervan.
- Tot slot werd dit rapport met meerderheid van stemmen goedgekeurd door de Raad van Bestuur.
- Alleen het KCE is verantwoordelijk voor de eventuele resterende vergissingen of onvolledigheden alsook voor de aanbevelingen aan de overheid.

Publicatiedatum: 14 januari 2013

Domein: Good Clinical Practice (GCP)

MeSH: Prostatic Neoplasms; Prostatectomy; Watchful Waiting

NLM classificatie: WJ 762

Taal: Nederlands, Engels

Formaat: Adobe® PDF™ (A4)

Wettelijk depot: D/2012/10.273/99

Copyright: De KCE-rapporten worden gepubliceerd onder de Licentie Creative Commons « by/nc/nd » <http://kce.fgov.be/nl/content/de-copyrights-van-de-kce-rapporten>.



Hoe refereren naar dit document?

Mambourg F, Jonckheer P, Piérart J, Van Brabant H. Nationale Praktijkrichtlijn voor de aanpak van gelokaliseerde prostaatcancer: deel 1. Good Clinical Practice (GCP). Brussel: Federaal Kenniscentrum voor de Gezondheidszorg (KCE). 2012. KCE Reports 194A. D/2012/10.273/99.

Dit document is beschikbaar op de website van het Federaal Kenniscentrum voor de Gezondheidszorg.



■ VOORWOORD

De internationale wetenschappelijke consensus is inmiddels overweldigend: het systematisch screenen van mannen op prostaatanker door middel van een PSA-test biedt voor deze mannen geen toegevoegde waarde, en verhoogt sterk het risico op een nutteloze ingreep, met alle gevolgen van dien: risico op incontinentie en impotentie. Deze conclusies zijn gebaseerd op state-of-the-art epidemiologisch onderzoek. En toch ... nog steeds blijven klinici – huisartsen en urologen – veel PSA-tests voorschrijven. Politieke figuren permitteren zich ongewettigde uitspraken over de weldaden van de agressieve aanpak – denken we maar aan de uitspraken van ex New-York burgemeester Rudolph G1uliani. Ook in ons land verschijnen tot op vandaag voorstellen, ja zelfs heuse provinciale initiatieven om georganiseerd te gaan screenen.

En dus vindt men – uiteraard – een verhoogde PSA-waarde bij een groot aantal mannen van middelbare en oudere leeftijd. En dus gaat men biopsies nemen. En dus vindt men regelmatig ook kwaadaardige cellen. De meeste van deze mannen hebben een kleine, vrij gelokaliseerde kanker, die eigenlijk niet (onmiddellijk) moet behandeld worden. Het gevolg is dat men hen opzadelt met een probleem waarmee ze natuurlijk meestal geen blijf weten. In se komt het hierop neer: “Meneer, u heeft kanker, maar in feite is het niet echt nodig om te behandelen. Aan u de keuze”.

In de praktijk gaat het er trouwens vaak nog anders aan toe, en stelt men meteen een radicale prostaatoperatie voor. Uit de cijfers van de VS blijkt bovendien dat de opkomst van de chirurgische robotten alleen maar geleid heeft tot een enorme toename van PSA-tests en van prostaatoperaties, waarvan zeer vele geen voordeel bijbrengen voor de patiënt.

Voor deze groep mannen met een meestal “toevallig” gevonden prostaatanker werd deze richtlijn ontwikkeld. Wij willen aan hun artsen een richtlijn aanreiken over de huidige wetenschappelijk verantwoorde aanpak van dit probleem. Het zal nooit een gemakkelijke beslissing of boodschap zijn, en daarom zullen wij ook een studie wijden aan de attitudes en barrières bij de betrokken actoren. Ten slotte volgt later ook nog een praktijkrichtlijn voor de aanpak van de meer geavanceerde prostaatankers.

Gewoontegetrouw werd deze kankerrichtlijn ontwikkeld samen met het College voor Oncologie; wij danken hen voor deze gewaardeerde samenwerking

Raf MERTENS
Algemeen Directeur



■ SAMENVATTING

INLEIDING

Dit rapport maakt deel uit van een meer uitgebreid project dat als doel heeft klinische praktijkrichtlijnen met betrekking tot prostaatanker op te stellen. In dit eerste rapport wordt ingezoomd op “Watchful Waiting” (oplettend afwachten) en “Active Surveillance” (actieve opvolging) bij de behandeling van gelokaliseerde prostaatanker. De meeste van deze patiënten zijn asymptomatisch. De kanker werd veelal ontdekt naar aanleiding van een PSA-test. Bij sommigen werd hij toevallig gevonden omdat een endoscopische prostaatrectomie uitgevoerd werd voor een goedaardige prostaathypertrofie. Alle aanbevelingen die in dit rapport geformuleerd worden zijn gebaseerd op klinische werkzaamheid. De kosteneffectiviteit werd niet geanalyseerd. Het is de bedoeling dat onderhavige aanbeveling opgevolgd wordt door alle zorgverleners betrokken bij de behandeling van patiënten nadat de diagnose van gelokaliseerde prostaatanker gesteld werd.

DEFINITIES

Gelokaliseerde prostaatanker wordt gedefinieerd als een kanker die niet verder reikt dan het prostaatkapsel (cT1a-T2c N0M0). Hij wordt volgens zijn risico op progressie onderverdeeld in drie subcategorieën. Deze indeling wordt aangeduid als de d’Amico classificatie. Ze wordt volgens drie criteria opgesteld: het TNM stadium, de Gleason score die de agressiviteit van de kankercellen meet (graad van differentiatie) en de PSA-waarde. De risicocategorieën worden momenteel als volgt door de European Association of Urology (EAU) gedefinieerd:

- Laag risico (T1-2a en Gleason<7 en PSA<10ng/ml)
- Matig risico (T2b-c of Gleason=7 of PSA tussen 10 en 20 ng/ml)
- Hoog risico (T3a of Gleason>7 of PSA>20ng/ml)
- **"Watchful Waiting" bestaat erin bij patiënten die niet in aanmerking komen voor een curatieve behandeling af te wachten met behandelen. Bij deze strategie worden de patiënten verder opgevolgd en wordt een palliatieve behandeling ingesteld wanneer er kankergerelateerde symptomen optreden.**



- **"Active Surveillance" bestaat erin de behandeling van patiënten die in aanmerking komen voor een onmiddellijke curatieve behandeling uit te stellen. Bij deze strategie wordt de evolutie van de kanker periodiek opnieuw geëvalueerd en wordt bij progressie ervan een curatieve behandeling ingesteld.**

De levensverwachting van Belgische mannen (op bevolkingsniveau) is afkomstig van de desbetreffende tabellen over de jaren 2007-2009. De individuele levensverwachting van een bepaalde patiënt dient te worden berekend met een gevalideerd instrument dat rekening houdt met coorbiditeiten. De richtlijnontwikkelaars hebben voorgesteld om de keuze van dit instrument over te laten aan de leden van het multidisciplinaire oncologisch consult (MOC).

ONDERZOEKSVRAGEN

Bij het opmaken van deze praktijkaanbeveling werd uitgegaan van de volgende vragen:

1. Watchful Waiting versus Active Surveillance:
 - Welke patiënten kunnen baat hebben bij een Watchful Waiting of een Active Surveillance strategie?
 - Verschillen de gevolgen op het vlak van mortaliteit, morbiditeit of levenskwaliteit bij een Watchful Waiting of een Active Surveillance strategie met die van een onmiddellijke curatieve behandeling van een gelokaliseerde prostaatanker?
2. Active Surveillance:
 - Leidt een verschil in het gevolgde Active Surveillance protocol tot een verschillend resultaat qua mortaliteit, morbiditeit of levenskwaliteit?
 - Hoe wordt Active Surveillance in de praktijk uitgevoerd?
 - Wanneer moet overgestapt worden van Active Surveillance naar onmiddellijke curatieve behandeling?

METHODOLOGIE

Bij het opmaken van deze praktijkrichtlijn werd gebruik gemaakt van de ADAPTE methodologie waarbij (inter)nationale richtlijnen aangepast worden aan de Belgische context. Er werd gezocht naar (inter)nationale aanbevelingen in Medline (OVID), EMBASE, National Guideline Clearinghouse, websites van richtlijnproducenten en websites van oncologische organisaties. De 20 weerhouden aanbevelingen werden geëvalueerd op hun kwaliteit door twee onafhankelijke onderzoekers, gebruik makend van het AGREE instrument.

Vervolgens werd voor elk van de klinische vragen een actualisering van de geselecteerde aanbevelingen doorgevoerd, uitgaande van aanvullende gegevens uit de databases Medline (OVID), EMBASE en Cochrane Database of Systematic Reviews. Onze meest recente literatuur onderzoek zoektocht dateert van mei 2012.

De eindaanbevelingen werden op basis van het aldus gevonden bewijsmateriaal uitgewerkt door een multidisciplinair team (onder wie de auteurs van onderhavig document). Aan elke aanbeveling werd een aanbevelingskracht en een bewijsniveau toegekend aan de hand van het GRADE systeem (Tabel 1 en 2). Deze aanbevelingen werden nadien voorgelegd aan andere experts (de stakeholders), waaronder vertegenwoordigers van verschillende artsorganisaties. Zij hebben de aanbevelingen van een score voorzien en tijdens de vergadering gediscuteerd. Aan de hand van een formele procedure werden deze aanbevelingen door externe deskundigen gevalideerd. Er werd akte genomen van mogelijke belangenconflicten.

**Tabel 1 – Niveaus van bewijskracht volgens het GRADE-systeem.**

Niveau van bewijskracht	Definitie	Methodologische kwaliteit van ondersteunend bewijsmateriaal
Hoog	We betrouwen er sterk op dat het werkelijke effect dicht bij het geschatte effect ligt	RCT's zonder ernstige beperkingen of overweldigend bewijs uit observationele studies
Matig	We hebben een matig vertrouwen in het geschatte effect: het werkelijke effect zal waarschijnlijk dicht bij het geschatte effect liggen, maar de mogelijkheid bestaat dat er een aanzienlijk verschil is	RCT's met ernstige beperkingen (inconsistente resultaten, methodologische beperkingen, indirect, of onnauwkeurig) of uitzonderlijk sterk bewijs uit observationele studies
Laag	Ons vertrouwen in het geschatte effect is beperkt: het werkelijke effect kan aanzienlijk verschillen van het geschatte effect	RCT's met zeer ernstige beperkingen of observationele studies of patiëntenreeksen
Zeer laag	We hebben erg weinig vertrouwen in het geschatte effect: het werkelijke effect zal waarschijnlijk aanzienlijk verschillen van het geschatte effect	

Tabel 2 – Graad van aanbeveling volgens het GRADE-systeem.

Graad van aanbeveling	Definitie
Sterk	De gewenste effecten van een interventie wegen duidelijk op tegen de ongewenste effecten (<i>de interventie moet in de praktijk gebracht worden</i>), of de ongewenste effecten van een interventie wegen duidelijk op tegen de gewenste effecten (<i>de interventie moet niet in de praktijk gebracht worden</i>)
Zwak	De gewenste effecten van een interventie wegen waarschijnlijk op ten opzichte van de ongewenste effecten (<i>de interventie moet waarschijnlijk in de praktijk gebracht worden</i>), of de ongewenste effecten van een interventie wegen waarschijnlijk op ten opzichte van de gewenste effecten (<i>de interventie moet waarschijnlijk niet in de praktijk gebracht worden</i>)

Een aanbeveling wordt hieronder met « GKP » (Goede Klinische Praktijk) gekwalificeerd wanneer er geen gegevens uit gerandomiseerde klinische studies over bestaan, maar er wel een consensus over is binnen het hogervermelde multidisciplinair team.



KLINISCHE PRAKTIJKAANBEVELINGEN

Aanpak

Evaluatie

Goede Klinische Praktijk

Voorafgaand aan het nemen van een therapeutische beslissing moet een evaluatie gebeuren in het kader van het multidisciplinaire oncologisch consult (MOC), rekening houdende met:

- de algemene gezondheidstoestand van de patiënt, zijn levensverwachting en zijn comorbiditeit
- de kwaliteit van de biopsie, de kenmerken en de risicocategorie van de tumor.

Informatie

Goede Klinische Praktijk

Patiënten die in aanmerking komen en ook kiezen voor een curatieve behandeling, moeten worden geïnformeerd over de verschillende algemeen aanvaarde aanpakmogelijkheden, rekening houdende met hun gezondheidstoestand, hun individuele levensverwachting en de risicocategorie van de tumor. Deze aanpakmogelijkheden omvatten minstens de active surveillance, (externe en interstitiële) radiotherapie en radicale prostatectomie. De mogelijke voor- en nadelen van de verschillende opties zou moeten met de patiënt worden besproken en toegelicht.

Patiënten met gelokaliseerde prostaatkanker en een levensverwachting van <10 jaar

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Watchful Waiting is de aanbevolen aanpak bij patiënten met een levensverwachting van <10 jaar of bij patiënten met ernstige comorbiditeit.	Sterk	Matig

Patiënten met gelokaliseerde prostaatkanker met laag risico die in aanmerking komen en kiezen voor een curatieve behandeling.

Aanbevelingen	Graad van aanbeveling	Niveau van bewijskracht
Een Active Surveillance zou moeten overwogen worden, rekening houdende met de persoonlijke voorkeur van de betrokken patiënt en de functionele status van zijn urinaire, seksuele en digestieve functies.	Sterk	Zwak
Dergelijke patiënten moeten worden geïnformeerd over het feit dat werd aangetoond dat een onmiddellijke behandeling binnen een tijdsspanne van 10 à 12 jaar geen enkel voordeel biedt ten aanzien van observatie.	Sterk	Matig



Patiënten met gelokaliseerde prostaatanker met matig risico die in aanmerking komen en kiezen voor een curatieve behandeling

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Omwille van de grote anatomopathologische variabiliteit van prostaatankers met matig risico, kan er geen algemene aanbeveling geformuleerd worden betreffende de Active Surveillance bij patiënten die behoren tot deze subgroep.	Zwak	Zwak

Patiënten met gelokaliseerde prostaatanker met hoog risico

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Een Active Surveillance is voor dergelijke patiënten niet aanbevolen.	Sterk	Zwak

Opvolging van actieve surveillance

Biopsienamen gedurende het jaar dat volgt op de diagnose

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Er wordt aanbevolen om niet later dan 1 jaar na de diagnose een nieuwe biopsie te verrichten.	Sterk	Zwak

Onderzoeken

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
De volgende opties kunnen overwogen worden tijdens follow-up: tweemaal per jaar, een PSA-dosering en een klinische controle; jaarlijks een of meerdere beeldvormende onderzoeken.	Zwak	Zwak

Routinebiopsienames

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Volgend op de biopsienamen die gebeurde ten laatste 1 jaar na de diagnose, is het aanbevolen om biopsienames te herhalen. De optimale timing hiervan is momenteel niet gekend.	Sterk	Zwak



Patiënten wiens persoonlijke levensverwachting tijdens de opvolgingsperiode daalt tot <10 jaar

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Aanbevolen wordt om een Active Surveillance te stoppen bij patiënten van wie de individuele levensverwachting tijdens de opvolgingsperiode daalt tot <10 jaar, bij patiënten die 80 jaar worden of die een ernstige comorbiditeit ontwikkelen. Een Watchful Waiting kan dan voorgesteld worden.	Sterk	Matig

Progressie van de kanker

Aanbeveling	Graad van aanbeveling	Niveau van bewijskracht
Elke progressie van de kanker die wordt vermoed aan de hand van een PSA-waarde van >10ng/ml, een PSADT<3 jaar, een klinische verandering of een verdacht letsel bij beeldvorming, zou met een nieuwe biopsie moeten bevestigd worden en gevolgd worden door een herevaluatie van het risicocategorie.	Sterk	Zwak


Goede Klinische Praktijk

Indien een verhoging van risicocategorie bevestigd wordt, het zou moeten overwogen worden om van Active Surveillance over te schakelen naar onmiddellijke curatieve behandeling.



FLOWCHART

Aanpak






Levensverwachting van >10 jaar





Opvolging van Active surveillance



Indien een verhoging van risicocategorie bevestigd wordt, moet worden overwogen om van Active Surveillance over te schakelen naar onmiddellijke curatieve behandeling.

Aanbevolen wordt om een Active Surveillance te stoppen bij patiënten wier individuele levensverwachting tijdens de opvolgingsperiode daalt tot <10 jaar, bij patiënten die 80 jaar worden, of die een ernstige comorbiditeit ontwikkelen. Een Watchful Waiting kan dan voorgesteld worden.



IMPLEMENTEREN, EVALUEREN EN UPDATEN

Implementatie

Het implementeren van onderhavige aanbevelingen rond goede klinische praktijkvoering wordt gestuurd door het Nationaal College van Geneesheren Oncologie. Er wordt een on-line implementeringsinstrument, vergelijkbaar met de begeleidende instrumenten bij eerdere aanbevelingen, uitgewerkt.

Bijwerking van de aanbevelingen

Wanneer er nieuwe essentiële gegevens beschikbaar komen worden deze meegedeeld op de website van het Nationaal College van Geneesheren Oncologie en moet er overwogen worden een update van de desbetreffende elementen in het huidige rapport uit te voeren.

Een herevaluatie van onderhavige praktijkrichtlijn moet elke 5 jaar gebeuren.

VERDER ONDERZOEK

Kwaliteit van de biopsie

Er is nood aan het opstellen van een richtlijn waarin de kwaliteitscriteria voor prostaatbiopten worden gedefinieerd (zowel voor biopten die worden afgenomen via echografie dan wel via nucleaire magnetische resonantie). (GKP)

Standpunt van de patiënten en van de artsen

In een afzonderlijk rapport wordt een kwalitatieve analyse voorgesteld van de factoren die een impact hebben op de besluitvorming van zowel de urologen/radiotherapeuten als van de patiënten met betrekking tot de verschillende behandelingsopties voor gelokaliseerde prostaatkanker en inzonderheid Active Surveillance. Die analyse zal gebaseerd zijn op face-to-face interviews van artsen en patiënten.

Bovendien is het KCE momenteel bezig met de update van zijn rapport over prostaatkanker screening met PSA.



■ TABLE OF CONTENTS

LIST OF TABLES	4
LIST OF ABBREVIATIONS.....	5
■ SCIENTIFIC REPORT	7
1. INTRODUCTION	7
1.1. DEFINITIONS	7
1.2. GUIDELINE SCOPE.....	8
1.2.1. Stakeholders consultation	8
1.2.2. Results.....	9
2. METHODOLOGY	9
2.1. GENERAL APPROACH	9
2.2. CLINICAL QUESTIONS	10
2.3. LITERATURE SEARCH	10
2.3.1. Search strategy	10
2.3.2. Quality appraisal.....	10
2.4. DATA EXTRACTION AND SUMMARY	11
2.5. FORMULATION OF RECOMMENDATIONS	11
3. WATCHFUL WAITING VERSUS ACTIVE SURVEILLANCE	12
3.1. INTRODUCTION	12
3.2. CLINICAL QUESTIONS	13
3.3. SELECTED STUDIES	13
3.3.1. Guidelines.....	13
3.3.2. Systematic Reviews	13
3.3.3. Randomised controlled trials	14
3.3.4. Unpublished randomised controlled trials	15
3.4. CLINICAL EVIDENCE	16
3.4.1. Mortality	16



- 3.4.2. Morbidity 17
- 3.4.3. Quality of life..... 18
- 3.5. GUIDELINES RECOMMENDATIONS 18
 - 3.5.1. Patients' point of view..... 18
 - 3.5.2. Evolution of concepts 19
 - 3.5.3. Risk assessment 19
 - 3.5.4. Treatment 19
- 3.6. DISCUSSION..... 22
- 3.7. GDG ASSESSMENT 23
- 3.8. RECOMMENDATIONS 25
 - 3.8.1. Quality of the biopsy 25
 - 3.8.2. First assessment 25
 - 3.8.3. Information..... 25
 - 3.8.4. Men with life expectancy < 10 years 25
 - 3.8.5. Low-risk localised prostate cancer 25
 - 3.8.6. Intermediate-risk localised prostate cancer..... 25
 - 3.8.7. High-risk localised prostate cancer 25
- 4. DESCRIPTION OF AN ACTIVE SURVEILLANCE MANAGEMENT 26**
 - 4.1. INTRODUCTION 26
 - 4.2. CLINICAL QUESTIONS 26
 - 4.3. STUDIES SELECTED 26
 - 4.3.1. First question 26
 - 4.3.2. Second question..... 26
 - 4.3.3. Third question..... 28
 - 4.4. DISCUSSION..... 30
 - 4.5. GDG ASSESSMENT 30
 - 4.6. RECOMMENDATIONS 31
 - 4.6.1. Biopsy one year after the diagnosis 31
 - 4.6.2. Other tests 31



4.6.3.	Routine biopsy.....	31
4.6.4.	Life expectancy < 10y.....	31
4.6.5.	Disease reclassification.....	31
■	APPENDICES.....	32
APPENDIX 1.	DEFINITIONS.....	32
APPENDIX 1.1.	PROSTATE CANCER STAGING.....	32
APPENDIX 1.2.	GLEASON SCORE.....	33
APPENDIX 1.3.	RISK STRATIFICATION.....	33
APPENDIX 2.	FIRST STAKEHOLDERS CONSULTATION.....	34
APPENDIX 2.1.	WEBSURVEY “PROSTATE” SHORT LIST STATEMENTS.....	34
APPENDIX 2.2.	WEBSURVEY “PROSTATE” RESULTS.....	36
APPENDIX 3.	LITERATURE SEARCH.....	42
APPENDIX 3.1.	CLINICAL PRACTICE GUIDELINES.....	42
APPENDIX 3.2.	SYSTEMATIC REVIEWS AND PRIMARY STUDIES FOR QUESTION 1.....	46
APPENDIX 3.3.	RANDOMISED CONTROL TRIALS FOR QUESTION 3.....	49
APPENDIX 4.	QUALITY APPRAISAL.....	51
APPENDIX 4.1.	GUIDELINES.....	51
APPENDIX 4.2.	SYSTEMATIC REVIEWS.....	52
APPENDIX 4.3.	RANDOMISED CONTROLLED TRIALS.....	53
APPENDIX 5.	DATA EXTRACTION TABLES.....	54
APPENDIX 5.1.	SYSTEMATIC REVIEWS.....	54
APPENDIX 5.2.	PRIMARY STUDIES FOR TREATMENT.....	58
APPENDIX 5.3.	PUBLICATIONS NOT INCLUDED.....	62
APPENDIX 5.4.	GUIDELINES.....	63
APPENDIX 6.	GRADE.....	67
APPENDIX 6.1.	LEVELS OF EVIDENCE.....	67
APPENDIX 6.2.	DOWN- OR UPGRADING THE EVIDENCE.....	67
APPENDIX 6.3.	STRENGTH OF RECOMMENDATIONS.....	67
APPENDIX 6.5.	GRADE PROFILE.....	68



APPENDIX 7.	LIFE EXPECTANCY TABLE IN BELGIUM	71
APPENDIX 8.	QUALITY CRITERIA FOR PROSTATE BIOPSY	73
APPENDIX 9.	EXPERTS AND STAKEHOLDERS FINAL REVIEW.....	74
■	REFERENCES	77

LIST OF TABLES

Table 1 – Initials definitions of watchful waiting and active surveillance	13
Table 2 – Target population for AS/WW in localised prostate cancer according to guidelines	21
Table 3 – Type and frequency of exams included in active surveillance protocols from clinical guidelines.....	27
Table 4 – Rapid appraisal based on Questions 7, 8, 10 of AGREE (version May 2009 and scale 1 to 7).....	45
Table 5 – Guidelines appraisal followig AGREE tool.....	51



LIST OF ABBREVIATIONS

ABBREVIATION	DEFINITION
AGREE	Appraisal of guidelines for research and evaluation
AHRQ	Agency for Healthcare Research and Quality
AS	Active surveillance
ASA	American Society of Anesthesiology
ASAP	Atypical glands suspicious for cancer
AUA	American Urological Association
CPG	Clinical practice guideline
CDSR	Cochrane Database of Systematic Review
DRE	Digital rectal examination
EAU	European Association of Urology
EBRT	External beam radiation therapy
ESUR	European Society of Urogenital Radiology
GDG	Guideline Development Group
GIN	Guidelines International Network
HGPIN	High-grade prostatic intraepithelial neoplasia
HR	Hazard ratio
HRQoL	Health-Related Quality of Life
ISUP	International Society of Urological Pathology
mpMRI	Multi-parametric Magnetic Resonance Imaging
M-A	Meta-analysis
MeSH	Medical Subject Headings
MRI	Magnetic Resonance Imaging
NICE	National Institute for Health and Clinical Excellence
NIH	National Institute of Health
NNT	Number needed to treat
OMAR	Office of Medical Applications of Research
PCOS	Prostate Cancer Outcomes Study
PIVOT	Prostate Cancer Intervention versus Observation Trial



PSA	Prostate specific antigen
RCT	Randomised controlled trial
RP	Radical prostatectomy
RT	Radiotherapy
RR	Relative Risk
SPCG4	Scandinavian Prostate Cancer Group Study 4
SR	Systematic review
TRUS	Transrectal Ultrasonography
TURP	Transureteral resection of prostate
VACURG	Veterans Administration Cooperative Urological Research Group
VIKC	Vereniging Integrale Kankercentra
WW	Watchful waiting



■ SCIENTIFIC REPORT

1. INTRODUCTION

1.1. Definitions

Localised prostate cancer (cT1a-T2c N0 M0) refers to the clinical condition where a cancer is confined to the prostate gland, in the absence of lymph node invasion or metastases, corresponding to a stage NOM0. Prostate cancer (T3a-T4 N0 M0) and any T N+ M0 are considered as loco-regionally advanced prostate cancer.

Gleason score and TNM classification for prostate cancer are shown in Appendix 1.1. Gleason score is the most commonly used system for grading adenocarcinoma of the prostate and is related to the degree of differentiation of the cancer tissue. It can only be assessed using biopsy material. The Gleason score is the sum of the two most common patterns (grades 1-5) of tumour growth found. The Gleason score ranges between 2 and 10, with 10 representing the most aggressive cancers. The correlation between the Gleason score and survival data was first assessed in the VACURG trial.¹

Localised prostate cancers were first classified by d'Amico based on TNM classification, Gleason score and PSA level (for more details, Appendix 1.3).

Localised prostate cancers (cT1a-T2c N0 M0) are actually classified into 3 categories according to the risk of progression by the European Association of Urologist.²

1. Low risk: T1-2a and Gleason < 7 and PSA < 10 ng/ml. This group also contains a subgroup of patients that is presently recognized as "indolent" disease, or very low risk. This includes patients with maximum 2-3 positives biopsy's core from a 12 cores biopsy, each of the positive cores containing maximum 20 to 50% of cancer.
2. Intermediate risk: T2b-c or Gleason 7 or PSA 10-20 ng/ml.
3. High risk: T3a or Gleason > 7 or PSA > 20 ng/ml.



Epidemiology

According to the Belgian Cancer Registry, 8.681 new cases of prostate cancer were diagnosed in Belgium in 2009 (< 60y: 1 354, > 60 and < 70y: 2 957, > 70 and < 80y: 3 110, > 80y: 1 260). As in European country, prostate cancer is the most frequent cancer in men in Belgium. Although the yearly number of diagnosed cases increases, the cumulative mortality of prostate cancer remains stable at around 1.1% for men younger than 75 years and at 3.3% for men older than 75 years.³

1.2. Guideline scope

This study aims to develop a clinical practice guideline (CPG) on the treatment of localised prostate cancer. Importantly, this CPG will not address screening for prostate cancer.

We decided which clinical questions to consider in this guideline after a stakeholders consultation (see point 1.3.1). This guideline will be developed in collaboration with the College of Physicians for Oncology and is intended to be used by all health care providers involved in the care for prostate cancer patients. It was decided to develop a number of consecutive separate reports, each discussing a limited number of clinical questions and recommendations.

1.2.1. Stakeholders consultation

Stakeholders were consulted in 2011. An inventory of relevant clinical questions was built in 4 steps, with the aim of identifying the most problematic issues:

- Step 1: The coordinator of the guideline development group (GDG) of the Oncology College of the Federal Public Service of Public Health sent to KCE experts a list of 28 questions about the following aspects of prostate cancer treatments: active surveillance, radical prostatectomy, external or internal (brachytherapy), beam radiation therapy (EBRT), hormone therapy, bone targeted therapies and chemotherapy. The KCE experts, in concertation with the coordinator then shortened this list to 11 questions identified as the most relevant ones in current Belgian practice.

- Step 2: KCE experts performed a rapid appraisal of existing guidelines and chose the methodological best rating guideline on the AGREE scale (<http://www.agreecollaboration.org>): the NICE Full Guideline Prostate cancer: diagnosis and treatment (80 recommendations).
- Step 3: The list send by Oncology College and the NICE recommendations were merged and shortened by KCE experts taking into consideration the scope of the guideline, i.e. localised prostate cancer, this ultimately lead to 19 potentially problematic recommendations.
- Step 4: To further limit the number of clinical questions to be scrutinised, the inventory of the 19 potentially problematic recommendations was presented by means of a websurvey, to practitioners involved in prostate cancer care and to patients. Surveyed practitioners were urologists, radiotherapists, general practitioners and nurses in urology. Surveyed patients were members of one Belgian patient association : "Wij ook" and one European association: Europa Uomo. These practitioners and patients were invited by their representatives (5 professional unions, 2 scientific societies and 2 patient groups) to indicate if they agreed or disagreed with the content of each of these 19 potentially problematic recommendations (see the letter and the questionnaire in appendices). Only two alternatives were presented (agree or disagree) to invite the respondents to mark clearly their preferences in one single round. This questionnaire was pre-tested amongst KCE experts (more details in Appendix 2).

Responses were analysed by KCE experts. The discrepancies between the participants' views were brought to the fore by choosing a threshold of 70% of agreement. Statements with a percentage of agreement inferior to this threshold were considered as problematic statements.



1.2.2. Results

Five themes emerged from the final list of recommendations:

- The role of watchful waiting/active surveillance in the management of localised diagnosed prostate cancer;
- Equivalence of treatment between surgery and radiotherapy in terms of efficacy and side effects;
- Equivalence of different modes of surgery in terms of efficacy and side effects: open surgery, standard laparoscopic surgery, robot-assisted laparoscopic surgery;
- Equivalence of different modes of radiotherapy (external or internal) in terms of efficacy and side effects;
- The place of hormone therapy in the management of localised prostate cancer in relation to operative risk profile, cancer stage and patient characteristics.

We choose to publish separately the part of the guideline that is focused on the role of watchful waiting/active surveillance in the management of localised prostate cancer.

2. METHODOLOGY

2.1. General approach

The present clinical practice guideline CPG was developed by adapting existing international CPGs to the Belgian context. This approach was recently structured in a formal methodology by the ADAPTE group, an international group of guideline developers and researchers.⁴

The ADAPTE methodology generally consists of three major phases (www.adapte.org):

1. Set-up Phase: Outlines the necessary tasks to be completed prior to beginning the adaptation process (e.g., identifying necessary skills and resources).
2. Adaptation Phase: Assists guideline developers in moving from selection of a topic to identification of specific clinical questions; searching for and retrieving guidelines; assessing the consistency of the evidence therein, their quality, currency, content and applicability; decision making around adaptation; and preparing the draft adapted guideline.
3. Finalisation Phase: Guides guideline developers through getting feedback on the document from stakeholders who will be impacted by the guideline, consulting with the source developers of guidelines used in the adaptation process, establishing a process for review and updating of the adapted guideline and the process of creating a final document.

If necessary, included guidelines were updated with more recent evidence.



2.2. Clinical questions

This part of the clinical practice guideline addresses the following clinical questions:

1. Who may benefit from watchful waiting or active surveillance? What are the comparative outcomes (in terms of mortality, morbidity and quality of life) of watchful waiting (WW) or active surveillance (AS) versus immediate treatment with curative intent for localised prostate cancer?
2. How is active surveillance (AS) organised?
 - Did different active surveillance strategies affect the outcomes?
 - How is active surveillance implemented?
 - When to switch from AS to curative intervention?

2.3. Literature search

2.3.1. Search strategy

The search for was performed step by step.

1. A first search of guidelines was done based on the comprehensive clinical questions.
2. An additional search for meta-analyses (M-A) and systematic reviews (SRs), to perform an update of the best quality guidelines, was done for each specific clinical question.
3. The search was extended to randomised control trials (RCTs) for question 1.
4. The search was extended to observational studies for the question 3 and will be described in this chapter.

To identify published CPGs on prostate cancer, Medline (through OVID), the National Guideline Clearinghouse and specific websites (see 3.1) were searched.

For Medline, the following MeSH or non MeSH terms were used in combination: "prostatic neoplasm" [MeSH Terms] OR prostate cancer [Text Word]. These MeSH terms were combined with a standardised search strategy to identify CPGs. Both national and international CPGs were searched. A language (English, Dutch and French) and date restriction

(>2000) were used. CPGs without references were excluded, as were CPGs without clear recommendations.

The search for meta-analysis (M-A) and systematic review (SR) included a search in Medline (through OVID), EMBASE and the Cochrane Database of Systematic Reviews (see Appendix 3.2 for search strings). As for the CPGs, the search was limited to articles published in English, French and Dutch. In general, systematic reviews not reporting the search strategy and/or the quality appraisal of the included studies were excluded. All searches were run between March 2011 and May 2012.

The identified studies were selected based on title and abstract. For all eligible studies, the full-text was retrieved. In case no full-text was available, the study was not taken into account for the final recommendations.

2.3.2. Quality appraisal

2.3.2.1. Clinical practice guidelines

After exclusion of duplicates, 20 guidelines were found on specific website. Two hundred and fifty six publications were retrieved on PubMed (Ovid). Based on title and abstract and after exclusion of duplicates, 17 publications issued from PubMed were selected. Guidelines not focused on watchful waiting and active surveillance and one Japanese guideline not relevant for our population were first excluded. We performed thereafter a rapid appraisal of guidelines quality based on questions 7, 8 and 10 included in AGREE tool.⁵ Those questions are focused on selecting the evidence. We consider a good evidence selecting process as a basic requirement and decide not to follow the appraisal if no good description of this process was available. Guidelines were finally scored using the full AGREE-2 tool by two independent researchers (FM and PJ) and discussed in case of disagreement (see Appendix 4.1 for an overview of the scores).

2.3.2.2. Systematic reviews

The quality of the retrieved SR and M-A was assessed using the checklists of the Dutch Cochrane Centre (www.cochrane.nl). All critical appraisals were done by a single KCE expert (see Appendix 4.2 for an overview of the scores).



2.3.2.3. *Randomised controlled trials*

The quality of the retrieved RCTs was assessed using the checklists of the Dutch Cochrane Centre (www.cochrane.nl). All critical appraisals were done by a single KCE expert (see Appendix 4.3 for an overview of the scores).

2.4. Data extraction and summary

For each included CPG the following data were extracted: search date and publication year, searched databases, availability of evidence tables, recommendations and referenced evidence.

For each systematic review, the search date, publication year, included studies and main results were extracted. For randomized controlled trials, the following data were extracted: publication year, study population, study intervention and outcomes.

For each clinical question, the recommendations were summarized in data extraction tables. Data extraction tables are provided in Appendix 5. If trial of sufficient quality was found, a level of evidence was assigned to each recommendation and additional study using the GRADE system (see Appendix 6).

2.5. Formulation of recommendations

Based on the retrieved evidence, a draft of recommendations was prepared by the KCE team (HV, FM, PJ). This draft together with the data extraction tables were circulated to the guideline development group (see Appendix 9) prior to each face-to-face meeting. The guideline development group met on 2 occasions (06/25/2012, 09/25/2012) to discuss the drafts. Recommendations were changed if important evidence supported this change. Based on the discussion meetings a second and a third draft of recommendations were prepared and sent for reviewing to the GDG members.

The recommendations prepared by the guideline development group were circulated to the Professional Associations (see stakeholders list). These panellists received the recommendations one week prior one open meeting (11/27/2012). As a preparation of the meeting all invited panellists were asked to score each recommendation on a 5-point Likert-scale to indicate their agreement with the recommendation, with a score of '1' indicating 'completely disagree', '2' indicating 'somewhat disagree', '3' indicating 'unsure', '4' indicating 'somewhat agree', and '5' indicating 'completely agree' (the panellists were also able to answer 'not applicable' in case they were not familiar with the underlying evidence). In case a panellist disagreed with the recommendation (score '1' or '2'), (s)he was asked to provide appropriate evidence. All scores (n = 15) were then anonymously summarized into a mean score, standard deviation and % of 'agree'-scores (score '4' and '5') to allow a targeted discussion (see Appendix 9).



3. WATCHFUL WAITING VERSUS ACTIVE SURVEILLANCE

3.1. Introduction

The long natural history of prostate cancer and the fact that many men diagnosed with prostate cancer may ultimately die from other causes can explain the dilemma one faces in making a choice between an immediate curative treatment and an observational approach with deferred treatment. The latter can be accomplished through two distinctive approaches:

- watchful waiting, which involves a policy of observation and the provision of (palliative) treatment when symptoms arrive; and
- active surveillance which involves close monitoring for biochemical or histological progression with initiation of curative therapy at a given moment.

Unfortunately, the terms watchful waiting and active surveillance are used inconsistently by different authors. Some of them use the terms interchangeably, whereas others draw a clear distinction between them.⁶ Original definitions of watchful waiting and active surveillance are summarised by Parker in Table 1.⁷



Table 1 – Initials definitions of watchful waiting and active surveillance

	Active Surveillance	Watchful Waiting
Primary Aim	To individualize treatment	To avoid treatment
Patient Characteristics	Fit for radical treatment; age 50-80	Age >70 or life expectancy <15 years
Tumor Characteristics	T1-T2, Gleason ≤7, Initial PSA <15	Any T stage, Gleason ≤7, Any PSA
Monitoring	Frequent PSA testing, Repeat biopsies	PSA testing unimportant, No repeat biopsies
Indications for Treatment	Short PSA doubling time, Upgrading on biopsy	Symptomatic progression
Treatment Timing	Early	Delayed
Treatment Intent	Curative	Palliative

3.2. Clinical questions

This chapter addresses two questions:

- What are the comparative outcomes (in terms of mortality, morbidity and quality of life) of watchful waiting (WW) and active surveillance (AS) versus immediate treatment with curative intent for localised prostate cancer?
- Who may benefit from watchful waiting or active surveillance?

3.3. Selected studies

3.3.1. Guidelines

We selected three guidelines⁸⁻¹⁰ of high quality and two^{11,2} of moderate quality according to AGREE score. Their recommendations are summarised below. All guidelines included the two RCTs described below.^{1,12}

3.3.2. Systematic Reviews

Two SRs of high quality were selected.^{6,13}

- Wilt published in 2008 a SR aiming to determine the comparative benefits and harms of therapies for clinically localised prostate cancer and how patients and tumour characteristics affect the outcomes of these therapies. This SR was funded by AHRQ and the last search date for RCTs and observational studies was done in September 2007. In this SR, watchful waiting encompasses expectant management or active surveillance. Only 2 randomised trials comparing effectiveness between WW and immediate treatment were selected by the author.^{1, 12} VACURG trial and SPCG-4 trial are described in point 3.3.3. None of these RCTs enrolled patients with prostate cancer that was primarily detected by PSA testing. Wilt used also 473 observational studies (>80% cases series). However we could not find how many of them were related to WW. Additional data on harms and patients satisfaction came from the The Prostate Cancer Outcomes Study (PCOS), a large, nationally representative prospective survey of men with localised prostate cancer diagnosed in 1994 or 1995.¹⁴



- Hegarty published in 2008 one Cochrane SR aiming to compare the beneficial and harmful effects of curative treatment versus WW for the treatment of localised prostate cancer. The systematic search for randomised or quasi-randomised control trials was conducted from 1966 to 30th July 2010. Hegarty retained the two randomised trials selected by Wilt but no observational studies.⁶

Two SRs of moderate quality were also selected.^{15, 16}

- The SR published in 2011 by Agency for Healthcare Research and Quality (AHRQ) focused on active surveillance in men with localised prostate cancer. This SR was funded by National Institutes of Health (NIH-US) Office of Medical Applications of Research (OMAR) and the last search date for RCT and observational studies was done in August 2011. This SR found no trial on the effectiveness of AS but 2 updates of VACURG and SPCG-4 trials and 16 cohort studies (mainly retrospective) for other observational management strategies (largely resembling WW). Single-arm AS cohort studies were not included because they could not address comparative effectiveness questions. Moreover, the authors outline the likelihood of confounding by indication in the included cohort studies, due to the differences in patient characteristics and risk profile between patients treated with observational strategies and those who received active treatments. This SR is appraised as of moderate quality because the literature search was focused on Medline and CDSR only, and it's no clear if the quality appraisal of the primary studies was taken into account. Nevertheless, this SR is used here as a basis of process description of AS.¹⁵
- The SR provided by Farmaka, graded of moderate quality, was also included. This SR was done in order to prepare a Belgian consensus conference on the treatment of a variety of prostate pathologies. One question of the search was: "In which cases, and based on what characteristics, is active surveillance justified as a management strategy in patients with localised prostate cancer?" This SR was funded by INAMI-RIZIV (Belgium) and the search date was August 2010¹⁶. This SR identified the 2 abovementioned RCT and several observational studies. The authors stressed the very low quality of evidence of the VACURG trial and quoted many limitations of the

observational comparative studies (e.g. no clear distinction between WW and AS, wide variety of surveillance protocol, selection of often younger and healthier men). Some cohort studies where the total population was monitored by active surveillance were retained despite their short follow-up period (maximum 7 years) and different active surveillance protocols.

3.3.3. *Randomised controlled trials*

Five RCTs were selected. SPCG-4 and VACURG compared radical prostatectomy with watchful waiting. Although those two studies were included in the SR, we choose to describe them separately because before the publication of the PIVOT trial, they represented the state of the art. Three trials study an active surveillance regimen: one compares expectant management (WW) versus radical prostatectomy (PIVOT) and two focus on active surveillance versus radical prostatectomy or radiotherapy (Protect, Start). Furthermore we found two trials with quality of life as primary outcome, one related to men participating in the UMEA trial and the other to men participating in the SPCG4 trial.¹⁷ The UMEA-1 trial compared the efficacy of external beam radiotherapy versus WW and was conducted between 1986 and 1996 but remains unpublished so far. A long term follow-up focusing on quality of life of the 72 surviving patients was performed in 2001 by Fransson.¹⁸ Several measures of the quality of life of the patients involved in the SPCG4 trial were also performed.¹⁷

All RCTs are briefly described below.

3.3.3.1. *VACURG trial*

The VACURG trial started in the pre-PSA screening era and was the first RCT comparing radical prostatectomy versus expectant treatment for early carcinoma of the prostate. This trial started in 1967 and was conducted by Iversen in 15 Veterans Administration Hospitals in U.S. This RCT is appraised as of poor quality because in the absence of bone scan and diagnostic lymphadenectomy in routine evaluation, it is plausible that patients with disseminated disease were enrolled. Furthermore, sample size was small (< 300 participants).¹



3.3.3.2. SPCG-4 trial

The SPCG-4 started in 1989 and was the second RCT comparing radical prostatectomy (RP) versus watchful waiting (WW). This trial was funded by the Swedish Cancer Society and the National Institutes of Health and was conducted in 14 centres in Sweden, Finland and Iceland.

This RCT of high quality enrolled 695 men at the beginning of the PSA screening era. Patients had newly diagnosed, localised prostate cancer T0d, T1, T2 (following criteria available in 1978). Only 12% of the patients had no palpable T1c tumours at the time of enrolment in the study. Tumours were well or moderately well differentiated and predominantly detected by symptoms, rather than PSA. Patients were relatively young (< 75 years old) with a life expectancy of more than 10 years. WW was strictly defined and included no initial treatment. In the intervention group, the surgery had a radical nature and was not primarily aimed at the preservation of sexual potency. Tumour progression in the watchful waiting group was defined as palpable extracapsular extension or symptoms of obstruction requiring intervention. Hormonal therapy was given if metastases were detected by bone scan or, after 2003, if signs of tumour progression, including elevations of PSA level appeared. Follow-up occurred every 6 months (clinical examination and blood test thereafter) for the first 2 years, then annually (including bone scan and chest x-ray annually before 2003, bone scan every 2 years after).¹⁹ The most recent data issued from SPCG-4 after a follow-up of 15 years were published in 2011 by Bill-Axelsson.¹²

3.3.3.3. PIVOT trial

The PIVOT trial is a multicenter RCT comparing radical prostatectomy versus observation for men with localised prostate cancer. This trial was supported by the Department of Veterans Affairs Cooperative Study Program, the US National Cancer Institute and the US Agency for Healthcare Research and Quality (AHRQ). The trial was conducted in 44 Veterans Administration Hospitals and 8 National Cancer Institute sites across the U.S. The primary outcome of this trial was all-cause mortality; secondary endpoints were prostate cancer mortality and the occurrence of bone metastases. This high quality RCT enrolled 731 men with localised prostate cancer from November 1994 through January 2002. They had to be less than 76 years of age with an expected life expectancy of more than

10 years, and judged to be medically and surgically fit for RP. They were randomly assigned to radical prostatectomy (RP) or observation and followed through January 2010. Patients included had new (diagnosed within the past 12 months) biopsy proven clinically localised prostate cancer (T1-T2, NxM0). In three quarters of the population, the primary reason for biopsy was a PSA elevation or rise. Their mean age was 67 years and the median PSA value was 7.8 ng/mL. Approximately half of tumours (48% in RP group and 49.9% in observation group) were not palpable at digital rectal examination. Follow-up occurred every 6 months (with PSA measurement) for a minimum of 8 years, a maximum 15 years or until death. Quality of life was assessed every year, and bone scan performed every 5 years. In the intervention group (n=364), the technique used for radical prostatectomy, as well as additional interventions were at the surgeon's discretion. In the observation group (n=367), interventions for asymptomatic progression (e.g. change in PSA value) were discouraged. Palliative (non-curative) therapy (TURP, AD and/or targeted RT) was generally reserved for symptomatic or metastatic disease progression.²⁰

3.3.4. Unpublished randomised controlled trials

3.3.4.1. ProtecT trial

The ProtecT trial is an ongoing multicenter three-arm RCT. It will assess the clinical effectiveness, cost-effectiveness, and acceptability of active monitoring, radical prostatectomy and radical radiotherapy for men with localised prostate cancer. This trial started in 2001 and is conducted in 9 clinical centres in the UK. Recruitment is now complete and over 2500 men with prostate cancer are taking part in the ProtecT trial. The average age of participants is 64 years (range 50 to 80 years). So far, two years of follow up have been completed of a follow-up period of 10-15 years planned. First results should be published after 2015.²¹



3.3.4.2. *Start trial*

The trial started in 2006 and was conducted in Canada, US, England and Europe. It aimed to compare active monitoring versus radical prostatectomy and (external or internal) radiotherapy for men newly diagnosed with low risk prostate cancer. Unfortunately, this study was stopped early due to poor recruitment.²² No interim report was published. We contacted the principal investigator of the trial (Chris Parker), who could not provide reasons underlying the poor recruitment

3.4. Clinical evidence

3.4.1. *Mortality*

3.4.1.1. *Watchful waiting*

Systematic reviews

- The effect on mortality of interventional treatment vs watchful waiting was based on the SPCG-4 trial in the 4 SRs. They reported that radical prostatectomy improved prostate cancer survival compared with watchful waiting after a median of 8.2 years of follow-up.²³ Meanwhile, more recent recent results from the SPCG-4 study have become available and will be discussed below.
- Two SR added observational studies in their analysis although they differed in their results:^{13, 15}
 - Wilt did not formulate a conclusion because of the wide variation across studies (with overlapping overall survival estimates within and between treatments).¹³
 - According to AHRQ, radiotherapy was associated with a lower all-cause mortality rate than watchful waiting.
 - The retrospective studies were also consistent with those of the RCTs and suggested that radical prostatectomy was associated with a lower prostate-cancer-specific mortality than watchful waiting.¹⁵

Randomised controlled trials

- The major results of the low-quality VACURG trial showed not significant difference in median survival.
- After a follow-up of 15 years, the SPCG-4 showed that:¹²
 - All cause mortality
Radical prostatectomy was associated with a reduction from all cause mortality: RR=0.75 (0.61 to 0.92).
According to a post hoc statistical sub-group analysis, the NNT to avert one death was 15 overall and 7 for men younger than 65 years of age.
 - Prostate cancer mortality
Radical prostatectomy was associated with a reduction in the rate of death from prostate cancer: RR=0.62 (0.44 to 0.87).
- After a median follow-up of 10 years, the PIVOT trial²⁴ showed that:
 - All cause mortality
Radical prostatectomy did not significantly reduce all-cause mortality: HR=0.88 (0.71 to 1.08); p=0.22.
According to a preplanned sub-group analysis:²⁰
Among men with low-risk tumours (n=296), radical prostatectomy increased not significantly all-cause mortality: HR=1.15 (0.80 to 1.66).
Among men with intermediate-risk tumours (n=249), radical prostatectomy reduced significantly all-cause mortality: HR=0.69 (0.49 to 0.98).
Among men with high-risk tumours (n=157), radical prostatectomy reduced not significantly all-cause mortality: HR=0.40 (0.16 to 1.00).
Among men with PSA > 10, radical prostatectomy reduced all-cause mortality: HR=0.67 (0.48 to 0.94); p=0.22.
 - Prostate cancer mortality
Radical prostatectomy did not significantly reduce prostate cancer mortality: HR=0.63 (0.36 to 1.09); p=0.09.



3.4.1.2. Active surveillance

Systematic reviews

- Each SR mentioned the lack of RCT comparing interventional treatment with active surveillance in localised prostate cancer.
- Two SR added observational studies to consider this issue.^{15, 16} Nevertheless they differed in their conclusion:
 - According to AHRQ, evidence was insufficient to evaluate the comparative effectiveness of AS management versus immediate definitive treatment in men with localised prostate cancer (AHRQ). Moreover, the authors underlined the need of a standard definition of AS that clearly distinguishes it from WW.¹⁵
 - On the basis of cohort studies, the authors of Farmaka concluded that active surveillance in men with early stage prostate cancer was associated with low specific and total mortality rates (maximum follow-up <7 years). Surveillance active was considered therefore as a possible option in the management of early prostate cancer. The exact place, the type of patients who could get the benefit of this approach and the most suitable monitoring protocol remained to be defined.¹⁶

3.4.2. Morbidity

3.4.2.1. Watchful waiting

Systematic reviews

- The effect of immediate treatment vs watchful waiting on morbidity was based on the SPCG-4 in the SR that considered this question.^{6, 13, 15} Based on this RCT, an increased risk for urethral stricture among patients after radical prostatectomy was mentioned. They also quoted an increased relative risk for sexual dysfunction after radical prostatectomy compared with watchful waiting (Hegarty, Wilt). Hegarty reported that after a follow-up of 4 years, radical prostatectomy appeared to increase the risks of erectile dysfunction (RR = 1.78 (1.48 to 2.15)) and urinary leakage (RR = 2.29 (1.63 to 3.22)) although confident statements cannot be made about how frequently these adverse effects occur.⁶

- Two SR added results of observational studies on the morbidity analysis:^{13, 15}
 - According to the Prostate Cancer Outcomes Study results, inability to attain an erection was higher in men undergoing active intervention, especially androgen deprivation (86%) or radical prostatectomy (58%), than in men receiving watchful waiting (33%).¹³
 - According to AHRQ, men treated with radiotherapy had a higher rates of urinary strictures compared with men on WW.¹⁵

Randomised controlled trials

- After a follow-up of 15 years, the SPCG-4 showed that:¹²
 - The relative risk of distant metastases was lower in the radical prostatectomy group versus the watchful waiting: RR=0.59 (0.45 to 0.79).
 - The relative risk of local progression was lower in the RP group than in the WW: RR=0.34 (0.20 to 0.45).
 - The 1-year cumulative incidence of postoperative complications showed that the most common symptom reported after RP was impotence (58% of patients), followed by urinary leakage (32%).
- After a median follow-up of 10 years, the PIVOT trial²⁴ showed that:
 - Radical prostatectomy reduced significantly bone metastases: HR=0.40 (0.22 to 0.70); p< 0.001.
 - The perioperative complications occurred in 21.4% of men (wound infection was the most common) and included one death.
 - The patient-reported urinary incontinence was significantly more common in the RP group (17%) than in the observation group (6.3%), p< 0.001.
 - The patient-reported erectile dysfunction was significantly more common in the RP group (81.1%) than in the observation group (44.1%), p< 0.001.

3.4.2.2. Active surveillance

Systematic review

The lack of RCT comparing intervention with active surveillance in localised prostate cancer was also mentioned by the SR broaching the effect on morbidity.^{6, 13, 15} Moreover, no observational studies reporting clinical outcomes related to AS as compared to immediate definitive treatment were identified by the authors. AHRQ argued that efficacy results from studies of WW may represent the lower bound of the potential efficacy of AS.

3.4.3. Quality of life

3.4.3.1. Watchful waiting

Systematic reviews

- Three SRs broached this question.^{6, 13, 15}
- Two SRs mentioned also observational studies:^{13, 15}
 - According to the Prostate Cancer Outcomes Study results, patient satisfaction with all selected treatments was high and more than 90% would make the same treatment decision again, and most were delighted or pleased with their treatment decision. However, satisfaction was higher in men who received early intervention than in those who received watchful waiting.¹³
 - No conclusions was formulated in the AHRQ review because the results of observational studies varied across different domains of quality measure for radical prostatectomy and radiotherapy.¹⁵

Quality of life trials

- In the evaluation done by Johansson during the SPCG-4 trial comparing radical prostatectomy with watchful waiting:
 - No difference was found in self-assessed quality of life after more than 10 years between the 2 groups: no difference in number of physical symptoms (RP-WW: 94% vs. 94%).
 - Men were distressed more often by their erectile dysfunction and by urinary leakage when assigned to radical prostatectomy than to watchful waiting.

For instance, the RR of distress due to a loss of erection was 1.30 (1.00 to 1.70) and the RR of distress due to urinary leakage was 3.79 (2.36 to 6.06) in the RP group vs the WW.

- Moreover, the results showed lower scores in all psychological measures in the 2 groups versus a background population. For example, moderate to high level of anxiety were reported by the same proportion of patients in RP (43%) and WW (43%) but by fewer men in the control group (RR for patient group = 1.42 (95% CI: 1.07 to 1.88)).¹⁷
- In the evaluation done by Fransson during the UMEA-1 trial comparing external beam radiotherapy with watchful waiting :
 - There were no statistically significant differences at 10 years for HR QoL nor urinary trouble (mean 4.8 vs 3.0; p=0.034) nor bowel symptoms.
 - Sexual trouble appear to be more frequent (mean: 7.4 vs. 3.8, p=0.011) after radiotherapy than after watchful waiting (although no difference was found in erectile function nor in maintaining a sufficient erection).²⁵

3.5. Guidelines recommendations

3.5.1. Patients' point of view

Patient information was a considerable issue for NICE, AUA and VIKC. All three guidelines wrote specific recommendation therefore (see extraction table in Appendix 5.4.1). This topic was not explicitly mentioned in the EAU guideline. This guideline recommended in fact deferred treatment for all patients not willing to accept side-effects of active treatment. Surprisingly, patients' point of view did nowhere appear in the Spanish guideline.¹¹

Furthermore, AUA and NICE underlined the role of patient preferences in treatment decision making: "no one treatment modality is preferable for all patients"¹⁰ and "making treatment decisions, taking into account the effects on quality of life as well as survival".⁹



3.5.2. Evolution of concepts

Historical definitions of watchful waiting and active surveillance were presented in point 3.1.

Watchful waiting was traditionally considered in elderly or less fit men to avoid any intervention as long as possible. It excluded radical treatment options.²⁶ NICE recommends now that men with localised prostate cancer who have chosen a watchful waiting regimen and who have evidence of significant disease progression (rapidly rising PSA level or bone pain) should be reviewed by a physician.⁹ It seems that watchful waiting strictly defined is now being replaced by deferred treatment, depending not only on clinical but on biological progression as well. Of note, some guidelines uses the concept of deferred treatment^{8, 26} for both curative as palliative treatments.

Conversely, we found no significant evolution of the concept of active surveillance, representing an option for men who are fit for radical treatment in the event of disease progression.⁹

3.5.3. Risk assessment

NICE underscores first that an elevated PSA level alone should not automatically lead to prostate biopsy. A decision to take a biopsy must be done taking into account patient preferences, PSA level, DRE findings, and personal risks factors (family history).⁹

After diagnosis, AUA underscores that the first assessment must be extended to the overall health status and the life expectancy of the patient. Specific prostate cancer risk must be assessed on the same time (NICE). Therefore, specific prostate cancer risk strata that are significantly associated with PSA recurrence and cancer-specific mortality were defined. They use PSA, Gleason score and tumour stage. They define prostate cancer as of low, intermediate or high risk. Because variations of this system exist, we use here guidelines specifics definitions in extenso.

3.5.4. Treatment

3.5.4.1. Watchful waiting

- The Spanish guideline considers that watchful waiting may be a alternative for low risk localised prostate cancer in patients with limited life-expectancy (< 10 years).¹¹
- AUA underlines also that in the SPCG4 trial, radical prostatectomy may be associated with a lower risk of cancer recurrence and cancer specific mortality than watchful waiting.¹⁰ Based on the same trial, EAU recommends deferred treatment **only** for asymptomatic patients with well- or moderately-differentiated cancer, localised or locally advanced prostate cancer and a short life expectancy as for patient with high PSA levels (PSA < 50 ng/mL and PSA doubling time >12 months) for whom cure is unlikely.²⁶

3.5.4.2. Active surveillance

- EAU 2012 recommends active surveillance for Stage T1b-T2b patients who are well informed and have well-differentiated prostate cancer and a life expectancy of 10-15 years. Cancers with the lowest risk of progression are: PSA ≤ 10 ng/ml, biopsy Gleason score ≤ 6, ≤ 2 positive biopsies, ≤ 50% cancer per biopsy, cT1-2a. For those patients, active surveillance is recommended with a re-evaluation with PSA, TRUS and biopsies.²
- VIKC recommends active surveillance for low risk localised (T1c-2a, Gleason < 7, PSA < 10 ng/mL) prostate cancer **only** for patients with limited life-expectancy (older than 75 years).⁸
- The Spanish guideline considers that active surveillance can be offered in patients with clinically localised prostate cancer at low risk, Gleason < 3 + 3, < 50% affected cylinders in the biopsy and PSA < 15 ng/ml. group.¹¹



- For AUA also, active surveillance as interstitial prostate brachytherapy, external beam radiotherapy, and radical prostatectomy are appropriate monotherapy treatment options for the patient with low-risk localised prostate cancer (PSA < 10 ng/mL and a Gleason score of 6 or less and clinical stage T1c or T2a).¹⁰
- NICE recommends that men with low-risk localised prostate cancer who are considered suitable for radical treatment should first be offered active surveillance. A subgroup of men for whom active surveillance is particularly suitable is defined as: men with low-risk localised prostate cancer who have clinical stage T1c, a Gleason score 3+3, a PSA density < 0.15 ng/ml/ml and who have cancer in less than 50% of their total number of biopsy cores with < 10mm of any core involved.⁹
-


Table 2 – Target population for AS/WW in localised prostate cancer according to guidelines

Watchful waiting or deferred treatment								
	Clinical stage	Gleason score	Serum PSA	Biopsy core	Life Expectancy	Age	Other conditions	Strength
Aragon	Clinically localised prostate cancer : cT1-cT2, N0-Nx, M0-Mx.				< 10 years			Moderate
EAU	T1b-T2b				10-15 years		Well informed patients	Low
EAU	T3-T4		PSA< 50ng/ml		short			Low
Active surveillance								
	Clinical stage	Gleason score	Serum PSA	Biopsy core	Life Expectancy	Age	Other conditions	Strength
EAU	T1a T1c-T2a	< 7 (≤ 6)	≤ 10 ng/ml	≤ 2 positive biopsies & ≤ 50% cancer in biopsy	< 10 years < 10 years		Patients who do not accept treatment-related complications	Moderate
VIKC	T1c-2a T3	< 7	< 10 ng/ml		< 10 years	>75 years	Or co morbidities	Low
Aragon	Clinically localised prostate cancer : cT1-cT2, N0-Nx, M0-Mx.	3+3	< 15 ng/ml	≤ 50% cancer in biopsy				Very low
AUA	T1c or T2a	< 7	≤ 10 ng/ml				Patients preferences; health conditions related to urinary, sexual, and bowel function	Low



AUA	T2b	7	>10< 20		
NICE	T1-T2a	< 7	< 10 ng/ml	Cancer in < 50% of total number of biopsy cores with < 10 mm of any core involved	Men suitable for radical treatment Low
	Particularly if T1c	3+3	< 0.15 ng/ml/ml		
NICE	T2b-T2c	7	>10< 20		Option

3.6. Discussion

Clearly, there is a wide variation in the recommendations formulated in the guidelines mentioned above. This may be explained by several mechanisms. As we have shown in this chapter, the body of solid evidence supporting the management of patients with localised prostate cancer is limited. The two oldest RCTs (VACURG and SPCG4) that compared a deferred with an immediate treatment included patients diagnosed before the PSA era. Those patients were symptomatic at the time of diagnosis whereas most actual patients are asymptomatic and diagnosed through a PSA test. The PIVOT trial was more appropriate because three quarters of patients enrolled were diagnosed after a PSA elevation or rise.²⁴ Unfortunately, this trial compared radical prostatectomy versus *observation*. In the observation group, interventions for asymptomatic progression (e.g. change in PSA value) were discouraged.

Before the publication of the PIVOT trial results, there were no results from RCTs that compared “watchful waiting/observation” with immediate intervention, guideline developers had to rely on observational studies. Obviously these are more prone to bias, which might explain why GDG made different recommendations based on the same studies.

Another explanation why recommendations differ across guidelines may be that guideline developers differ in the relative weight they attribute to the survival benefit obtained from a given strategy and its adverse effects. If a GDG considers prolongation of life as the primordial outcome, it may recommend an interventional strategy whereas the same study may lead another GDG to recommend a more conservative approach because side effects are considered unacceptable compared to a minor survival benefit. The degree to which patient preferences are taken into account may also be an element that explains the observed variations.



3.7. GDG assessment

A first draft of the report was discussed on the first GDG meeting (05/25/2012). The GDG underlines that the quality of the biopsy is key for defining the risk category of the prostate cancer. Nevertheless, the clinical pathway leading to the diagnosis of prostate cancer was not considered a priority in this practice guideline (see 1.2.2). The following requirements related to the quality of the prostate biopsy and based on a review made by one of the participating pathologist were accepted by GDG consensus:

- The role of the urologist or radiologist:
 - To provide adequate clinical information to the pathologist.
 - To provide adequate tissue samples for pathological examination (12 core biopsies are required, with at least a single prostatic gland per biopsy and average length of prostate tissue > 10 mm per biopsy).
 - To handle the biopsies in a way that will help the pathologist to identify and map cancer in the prostate (separate container for each biopsy is warranted).
- The role of the pathologist:
 - To provide accurate and concise reports using unequivocal terms. Categories of diagnoses should be limited to (1) Benign prostatic gland and stroma, (2) Inflammation, (3) High-grade prostatic intraepithelial neoplasia (HGPIN), (4) Atypical glands suspicious for cancer (ASAP) malignancy cannot definitely be excluded and (5) Prostate cancer.²⁷ The categories of diseases (3) and (4), i.e. HGPIN and ASAP should never be diagnosed as cancer and never treated as cancer by the clinician.
 - If prostate cancer is diagnosed, the pathology report should include: location and distribution of the tumour, histopathological type, extent of tumour involvement (number of involved cores, linear length of cancer in mm and/or percentage of cancer involvement of each core),²⁸ Gleason score including primary and secondary patterns (according to the 2005 ISUP Modified Gleason Grading²⁹), and if present perineural and/or lymphovascular invasion. The pathologist should refer to the reporting recommendations published by Fine SW et al. (2012) for

special Gleason grading scenarios such as the context of abundant high-grade cancer, prostate cancer variants, glomeruloid structures and/or the presence of a tertiary Gleason pattern in prostate biopsies.³⁰ Perineural and lymphovascular invasion are currently not considered as essential reporting elements for prostate needle biopsies by leading international urological pathologists (more details in Appendix 8).³⁰

Although transrectal-ultrasonography (TRUS)-guided biopsy became the accepted standard for PCa diagnosis in the 1990s,³¹ the GDG underlined that multiparametric magnetic resonance imaging (mpMRI) may have a role in the diagnosis of localised prostate cancer after a PSA elevation or rise. mpMRI currently includes T1- and T2-weighted images, dynamic contrast, diffusion weighting, and proton spectroscopy.³² In a recently published SR the accuracy of MRI-guided biopsy was compared to with standard TRUS-guided biopsy. Results from this meta-analysis pooled 16 discrete patient population (n=599) and found that MRI-guided biopsy detects clinically significant cancer in an equivalent number of men versus standard biopsy. This is achieved using fewer biopsies in fewer men. However, methodological limitations of the individual studies and varying definitions of endpoints restrict the external validity of this systematic review. The authors concluded that there is a need for a large study with clearly defined mpMRI criteria, standardised sampling, and a standard definition of clinical significance of a given tumour.³³

This gap in knowledge has as partly been field by the European Society of Urogenital Radiology (ESUR). It started a process to define recommendations on standardised methods for the detection, localisation, and characterisation of prostate cancer. Those were achieved by a consensus meeting of 16 experts originating from UK, BEL, FR, and NL³²,³⁴ and contained a unified scoring system for MRI named the Magnetic Resonance Prostate Imaging Reporting and Data System (MR PI-RADS). This scoring system has been validated in 2012 by a French prospective study. This study enrolled 129 men suspected for prostate cancer and referred for MRI after at least one set of negative biopsies. A threshold of ESUR-S ≥ 9 showed the following characteristics: sensitivity: 73.5%; specificity: 81.5%; positive predictive value: 38.2%; negative predictive value: 95.2%; and accuracy: 80.4%. In this situation, the ESUR scoring



system was shown to provide clinically relevant stratification of the risk of showing prostate cancer.³⁵

The abovementioned information represent a first start in conducting more robust clinical trials on the efficacy of mpMRI. Although one unpublished modelling study^a concluded that the use of MRI and MRI-guided biopsies appears to be an efficient strategy, more economic evaluations of this approach for prostate cancer diagnosis will be critical to determine whether this is an efficient approach for risk assessment in all or only selected men presenting with prostate cancer.³⁶ Results of those study will need to be incorporated in future updates of the present guidelines.

The GDC concludes that a guideline with criteria of a good quality biopsy (as for TRUS-guided as for MRI-guided biopsy) is needed not only for the pathologists but also for the urologists.

Next to cancer-related elements defining the risk category of a given patient, his life expectancy based on his general health status needs to be considered. This may be based on validated life expectancy scale. The GDG suggested leaving the selection of the scale to the discretion of the local team.

Furthermore, men's priorities, needs and concerns are highly related to their family and personal values and need to be considered along the entire disease management.

The guideline development group underlines that the currently most relevant sources of high quality evidence are SPCG4 and PIVOT. Unfortunately, both trials included a "watchful waiting/observation" management and not an active surveillance management. Hence, for the time being, one cannot make any conclusion on the long term effects of any kind of management on low risk non palpable prostate cancer, radical prostatectomy and radiotherapy included. This information should be discussed with the patient before any decision is taken. However, mortality data derived from "watchful waiting" or "observation" trials remain of

interest since one can reasonably expect mortality under "active surveillance" management not to exceed mortality observed in patients that were managed under a "watchful waiting/observation" regimen. PIVOT results showed that radical prostatectomy did neither significantly reduce all-cause nor prostate cancer mortality over 10 years as compared to "observation". Moreover, the patient-reported urinary incontinence and erectile dysfunction were significantly more common in the RP group than in the observation group.²⁴ Therefore, the risk/benefit balance of a radical treatment can be expected to be unfavourable for men with a life expectancy less than 10 years.

Definitions

- Watchful waiting consists of deferring treatment in patients with prostate cancer who are no candidate and/or suitable for immediate curative treatment. WW implies following up patients and only treating them with a palliative intent if symptoms appear.
- Active surveillance consists of deferring treatment in patients who are candidate and suitable for immediate curative radical treatment. AS implies revisiting periodically the status of the patient and treating upon progression, still with a curative intent.

Prostate cancer risk categories :

- o Low risk: T1-2a and Gleason < 7 and PSA < 10 ng/mL.
- o Intermediate risk: T2b-c or Gleason 7 or PSA >10< 20 ng/mL.
- o High risk: T3a or Gleason >7 or PSA >20 ng/mL.

^a The cost-effectiveness of MRI and MRI-guided biopsy versus TRUS-guided biopsy in the diagnosis of prostate cancer: a modelling study. Poster 2012. M.deRooij@rad.umcn.nl



3.8. Recommendations

3.8.1. Quality of the biopsy

A guideline with quality criteria of a prostate biopsy is needed not only for pathologists but also for urologists and radiologists (*good clinical practice - GCP*)

3.8.2. First assessment

Before any treatment decision can be made, a assessment should be undertaken including:

- the patient's overall health status, his individual life expectancy and comorbidities
- the quality of the biopsy and tumour characteristics (including the risk category) (*GCP*)

3.8.3. Information

A patient, eligible and opting for a strategy with curative intent, should be informed about commonly accepted initial managements with regards to his health status, individual life expectancy and tumour risk category. Commonly accepted initial managements include at least active surveillance, radiotherapy (external beam and interstitial), and radical prostatectomy. The estimated benefits and harms of each intervention should be explained and discussed with the patient. (*GCP*)

3.8.4. Men with life expectancy < 10 years

In patients with localised prostate cancer (all risk category) and individual life expectancy < 10 years or with important comorbidities watchful waiting with palliative intent is recommended. (*strong recommendation, moderate level of evidence*)

3.8.5. Low-risk localised prostate cancer

In patients with low-risk localised prostate cancer, eligible and opting for a strategy with curative intent, active surveillance should be considered as a management option, taking into account patient preferences and health conditions related to urinary, sexual, and bowel function. (*strong recommendation, low level of evidence*)

Men with low-risk localised prostate cancer must be informed that at the present time there is no demonstrated benefit within 10 to 12 years for immediate treatments as opposed to observation. (*strong recommendation, moderate level of evidence*)

3.8.6. Intermediate-risk localised prostate cancer

Because of the pathological heterogeneity of the patients with intermediate-risk localised prostate cancer, no general recommendation can currently be made on active surveillance in this subset of patients.

3.8.7. High-risk localised prostate cancer

In patients with high-risk localised prostate cancer, active surveillance is not recommended. (*weak recommendation, low level of evidence*)

4. DESCRIPTION OF AN ACTIVE SURVEILLANCE MANAGEMENT

4.1. Introduction

Based on the GDG's definition of "active surveillance" (see above), the aim of this chapter is to further define the notion of "reassessing periodically the status of the patient and treating with curative intent upon progression".

4.2. Clinical questions

This chapter addresses three questions related to the active surveillance (AS) strategy:

1. Do different active surveillance strategies affect outcomes?
2. How is active surveillance implemented?
3. When to switch from AS to another intervention with curative intent?

4.3. Studies selected

4.3.1. First question

In 2011, NICE performed an update of its Clinical Guideline⁹ in which it expressed concerns over variations in the way active surveillance is being used and performed. Consequently, a new topic on the most effective active surveillance protocol was introduced. The new clinical question was: "In men with prostate cancer followed by active surveillance, is the nature and repetition frequency of surveillance tests (PSA, DRE, MRI, biopsy...) associated with cancer specific survival, overall survival or the rate of radical intervention?" NICE found 14 studies relevant to this clinical question. Although several observational studies were found, no RCT comparing different active surveillance strategies was found in April 2011.³⁷ We performed an update of NICE's search in August 2012 (see Appendix 3.3). We found no studies comparing different active surveillance strategies. However, we identified 8 observational studies that we considered potentially relevant for questions 2 and 3.³⁸⁻⁴³

4.3.2. Second question

Only three guidelines provided descriptions of active surveillance protocols.⁹⁻¹¹ To obtain more information on the current practice, we studied the 8 abovementioned observational studies. After revision, we excluded case series including less than 400 men. We selected the PRIAS study (Europe),³⁸ the Johns Hopkins experience (US)⁴³ and the Canadian experience.⁴¹

The PRIAS study is an international observational prospective study conducted worldwide since December 2006. This study is facilitated by an electronic Web-based decision tool. Inclusion criteria according to protocol are: clinical stage T1c or T2; Gleason score is ≤ 6 , two or fewer biopsy cores invaded with PCa, PSA ≤ 10 ng/mL, PSA density ≤ 0.2 ng/mL. Over 2000 men have been included until now from over 100 participating centres in 17 countries. Risk reclassification on repeat biopsy during the first 4 years of follow-up has occurred in approximately 30% of biopsied men, although a switch towards active therapy has been performed in 22% of the total cohort.³⁸

One observational prospective cohort study is conducted at the Johns Hopkins Hospital since January 1995. Inclusion criteria are: very-low-risk cancers, defined by clinical stage (T1c), prostate-specific antigen density < 0.15 ng/mL, and prostate biopsy findings (Gleason score ≤ 6 , two or fewer cores with cancer, and $\leq 50\%$ cancer involvement of any core). A total of 769 men have been included until 2011. Curative intervention was recommended on disease reclassification on the basis of biopsy criteria. Overall, 255 men (33.2%) underwent intervention at a median of 2.2 years (range, 0.6 to 10.2 years) after diagnosis. From those, 188 men (73.7%) underwent intervention on the basis of disease reclassification on biopsy.⁴³

One prospective single arm cohort study founded by the Canadian Prostate Cancer Research Foundation is conducted in Toronto since November 1995. Inclusion criteria are: clinical stage T1c or T2a; Gleason score ≤ 6 , PSA ≤ 10 ng/mL. A PSA doubling time of less than 3 years was considered as an indication for intervention until 2009 and resulted in treatment in 14% of the cohort. Since 2009, a short doubling time has prompted an additional biopsy or mpMRI, and no longer serves as the sole reason for treatment. Upgrading (to Gleason 4+3 or greater) occurred in 8% of the cohort and was an indication for intervention, although,



upgrading to Gleason 3+4 was not. The proportion of patients remained on surveillance was 84, 72, and 62% at 2, 5, and 10 years. One additional result from this study is that for men aged 70 or more, the hazard ratio for non prostate to prostate cancer mortality is 33.3 (95% CI: 8.2 to 136).⁴¹

We summarised in table 3 below, the active surveillance protocols as described in selected guidelines and in table 4 (4.3.2), the active surveillance protocols as described in the selected observational studies.

Table 3 – Type and frequency of exams included in active surveillance protocols from clinical guidelines

	NICE		AUA		Spanish NHS	
	Investigations	Timing	Investigations	Timing	Investigations	Timing
History and Physical Exam	Physical examination ECOG Performance status DRE	Every 3 months for 2 years then every 6 months	Physical examination	Periodic	DRE	Every 3 months for 2 years then every 6 months
Biochemistry	Serum PSA	Every 3 months for 2 years then every 6 months	Serum PSA	Periodic	Serum PSA	Every 3 months for 2 years then every 6 months
Radiology (bone scan, chest x-rays, CTs)	As clinically indicated	-				
Other Investigations	TRUS guided biopsy (minimum of 8 cores) Tissue sample from serial needle biopsies for tissue banking	year 1, 4, 7, 10 and then every 5 years	Biopsy	Periodic	Biopsy(10 cylinders at least)	Year 1, 4, and 7

DRE (Digital Rectal Exam)



Table 4 - Type and frequency of exams included in active surveillance protocols from observational studies

	PRIAS		Johns Hopkins		Canada	
	Investigations	Timing	Investigations	Timing	Investigations	Timing
History and Physical Exam	Clinical examination	Every 6 months for 2 years, then every year	DRE	Every 6 months		
Biochemistry	Serum PSA	Every 3 months for 2 years then every 6 months	Serum PSA (total and free)	Every 6 months	Serum PSA	Every 3 months for 2 years then every 6 months
Radiology (bone scan, chest x-rays, CTs)	Bone scan whenever PSA>20	-				
Other Investigations	Biopsy	At year 1, 4, 7, 10	Biopsy(12 to 14 cores)	Every year	Biopsy	After 6-12 months, then every 3-4 years until 80y.

DRE (Digital Rectal Exam)

4.3.3. Third question

Below, we summarise criteria of disease progression described in four guidelines (table 5)^{2, 9-11} and in the three selected observational studies (table 6).^{38, 41, 43}


Table 5 – Criteria of disease progression in major practice guidelines

	NICE	AUA	EAU	Spanish NHS
PSA progression	a) PSA DT < 2 years, based on at least 3 separate measurements over a minimum of 6 months b) Final PSA > 8 ng/ml c) P value < 0.05 from a regression analysis of ln (PSA) on time		PSA doubling time with a cut-off value ranging between ≤ 2 and ≤ 4 years	PSA velocity > 1 ng/ml/year
Histological progression	Gleason pattern predominant 4 or + (i.e. Gleason 7 (4+3) or higher) in the re-biopsy specimen of the prostate performed at 18 months, 5 years, and 10 years, as per protocol	Increased grade, Increased stage	Gleason score progression ≥ 7 At re-biopsy (interval 1-4 years)	Higher degree or greater extension of tumour
Clinical progression	a) More than twice increase in the product of the maximum perpendicular diameters of the primary lesion as measured digitally b) Local progression of prostate cancer requiring TURP c) Development of ureteric obstruction d) Radiological and/or clinical evidence of distant metastasis	Increased volume		Evidence of locally advanced disease in DRE



Table 6 - Criteria of disease progression in observational studies

	PRIAS	Johns Hopkins	Canada
PSA progression	PSADT < 3 year after a year of inclusion in the study	Not used	PSADT < 3 year
Histological progression	Stage >3, Gleason score >6, > 2 biopsy core invaded	Gleason score >6, or > 2 biopsy core invaded, or >50% cancer involvement in any core	Higher Grade
Clinical progression		Reclassification by volume	If unequivocal palpable nodule, biopsy is done

Source: observational studies

4.4. Discussion

As we have seen in Chapter 3, the body of scientific evidence in support of one or another management strategy of patients with localised prostate cancer is limited. We found no randomised trials comparing outcomes according to different active surveillance strategies. Consequently, it is impossible to choose the most effective surveillance protocol based on solid data. The variation in inclusion criteria, protocols and patient populations in observational studies preclude a one to one comparison of surrogates and final outcomes.³⁸

The reliance on PSA kinetics as the sole reason for treatment is debatable. Loblaw investigated the proportion of patients who would have undergone treatment according to PSA triggers only. Based on the Canadian experience, he found that the proportion of patients who would have had a trigger for treatment ranged from 14% to 42% for histological or clinical progression, versus 37% to 50% for the PSADT triggers and 42% to 84% for the PSA velocity triggers.⁴⁴ Consequently, in the Canadian experience, PSA kinetics alone was no longer used as a driver for radical treatment.⁴¹ Surprisingly, a specific protocol adaptation in relation to a patient's age was found only in the Canadian experience. Its protocol included a biopsy every 3-4 years until the patient reached 80. Afterwards, biopsy is no longer performed.

The results of an active surveillance strategy are encouraging so far. However, prospective trials are needed to further optimise the inclusion and follow-up criteria (as the optimal biopsy timing) for active surveillance.³⁸

4.5. GDG assessment

This draft of the report was discussed on the second GDG meeting (09/25/2012). The GDG underlined that although the body of scientific evidence is limited, the role of the “confirmatory” biopsy is of crucial importance. This is in accordance with Bul who underlined that the short amount of time (median: 1.3 yr) from diagnosis to surgery in the current results of the PRIAS trial (n=2 079) is likely due to reclassification of risk because of understaging or undergrading at diagnosis.⁴⁵

The reliability of the PSA measurements was subject to a discussion. The analysis of the predictive ability of PSA kinetics performed on the prospective cohort conducted at the Johns Hopkins Hospital was cited. In patients included in this cohort who eventually underwent radical prostatectomy, nor PSA velocity (p=0.79) nor PSADT (p=0.87) were associated with the presence of unfavourable surgical pathology.⁴⁶ In one other American cohort study cited by the experts, the positive confirmatory biopsy was the only independent predictor of progression: HR 3.16, (95% CI: 1.41 to 7.09, p=0.005).⁴⁷ Although new prostate cancer biomarkers may be promising, they were not discussed because of lack of hard evidence so far.



Finally, a consensus was found in accordance to the conclusions of the Canadian experience and others cited above.^{41, 46, 47} A change in PSA level or in PSA kinetics may be considered as a trigger to perform more examinations including a biopsy. Nevertheless, disease reclassification and switch to radical treatment must be based on biopsy results.

According to the conclusions of the literature review reporting the absence of valid trials (RCTs) comparing different active surveillance strategies, each team can choose to adopt a protocol including routine biopsy performed at fixed time or a more flexible strategy adapted to each patient. So, a common biopsy protocol such the PRIAS protocol should be considered. Nevertheless, one more flexible protocol keeping in mind the quality of life of the patients, and the anxiety related to the biopsy may also be adopted.

The GDG recommended routine biopsy to be stopped in patients after reaching the age of 80 and in case a patient's life expectancy drops below 10 years. PIVOT results showed us that radical prostatectomy did not reduce prostate cancer mortality over 10 years as compared to "watchful waiting/observation" regimen. For older men (>70 y.), the risk to die from other cause is 33.3 higher than to die from prostate cancer.⁴¹ The mean life expectancy of a man aged of 80 years was 7.22 year in 2007 for Belgium (see Appendix 7).

4.6. Recommendations

4.6.1. Biopsy one year after the diagnosis

A repeat biopsy is recommended not later than one year after the diagnosis. (strong recommendation, low level of evidence)

4.6.2. Other tests

PSA measurements and clinical examination every six months can be considered. Imaging each year can be considered. (weak recommendation, low level of evidence)

4.6.3. Routine biopsy

After the biopsy performed within one year, repeat biopsies are recommended; there timing can currently not be defined. (strong recommendation, low level of evidence)

4.6.4. Life expectancy < 10y

In case of the individual life expectancy becomes < 10 year or after reaching the age of 80, or in case of the development of significant comorbidity, it is recommended to stop active surveillance and to offer watchful waiting with palliative intent. (strong recommendation, moderate level of evidence)

4.6.5. Disease reclassification

Disease progression as suggested by PSA>10ng/mL, or PSADT< 3 years, or clinical change, or suspicious lesions at imaging, should be confirmed by an additional biopsy and followed by risk reclassification. (strong recommendation, low level of evidence)

Switching to a radical treatment should be considered in case of risk reclassification. (GCP)



■ APPENDICES

APPENDIX 1. DEFINITIONS

Appendix 1.1. Prostate cancer staging

The TNM classification is used to stage prostate cancer. It describes the extent of the primary tumour (T stage), the absence or presence of spread to nearby lymph nodes (N stage) and the absence or presence of distant spread, or metastasis (M stage). Table 1 below describes TNM classification from Union for International Cancer Control (UICC - www.uicc.org). It represents the 7th edition (TNM-7) of the TNM classification that took effect from January 2010 on, and includes major modifications on PCA as compared to the 6th edition.

Prostate- 7th edition

<p>T1 Not palpable or visible</p> <p>T1a ≤5% or less</p> <p>T1b >5%</p> <p>T1c Detected by needle biopsy</p> <p>T2 Confined within prostate</p> <p>T2a ≤ half of one lobe</p> <p>T2b > half of one lobe</p> <p>T2c Both lobes</p> <p>T3 Through prostate capsule</p> <p>T3a Extracapsular</p> <p>T3b Seminal vesicle(s)</p> <p>T4 Fixed or invades adjacent structures</p> <p style="text-align: right;">No change from 6th</p>	<p>STAGE GROUPING (ANATOMIC)</p> <p style="text-align: right;"><u>(UICC)</u></p> <p>Stage I T1, T2a N0</p> <p>Stage II T2b-2c N0</p> <p>Stage III T3 N0</p> <p>Stage IV T4 N0</p> <p style="padding-left: 40px;">Any T N1</p> <p style="padding-left: 40px;">Any T Any N M1</p> <p style="text-align: right; color: orange;">Change from 6th Grade was in 6th</p>
---	---



Appendix 1.2. Gleason score

The Gleason score is the most commonly used system for grading adenocarcinoma of the prostate. The Gleason score can only be assessed using biopsy material (core biopsy or operative specimens). Cytological preparations cannot be used. The Gleason score is the sum of the two most common patterns (grades 1-5, depending on histologic differentiation) of tumour growth found. The Gleason score ranges between 2 and 10, with 2 being the least aggressive and 10 the most aggressive. In needle biopsy, it is recommended that the worst grade always should be included, even if it is present in < 5% of biopsy material.

Appendix 1.3. Risk stratification

Localized PCa (T1-T3a N0 M0) are usually separated into 3 categories according **to the risk of progression:**

Low risk: T1-2a and Gleason < 7 and PSA < 10 ng/ml. **This group also contains** a subgroup of patients that is presently recognized as “**indolent**” disease, or **very low risk**. This includes patients with maximum 2-3 positive biopsies core from a 12 cores biopsy, each of the positive cores containing maximum 20 to 50% of cancer.

Intermediate risk: T2b-c or Gleason 7 or PSA 10-20 ng/ml.

High risk: T3a or Gleason > 7 or PSA > 20 ng/ml.



APPENDIX 2. FIRST STAKEHOLDERS CONSULTATION

Appendix 2.1. Websurvey “prostate” short list statements

	Français	Nederlands
2. CTscanpelvis	Il n'est pas recommandé de réaliser un CT scan du pelvis en cas de cancer de la prostate localisé à risque faible ou intermédiaire.	Een CT-scan van het bekken is niet aanbevolen bij mannen met een laag- of intermediair risico gelokaliseerd prostaatCA.
3.Scintigraphier	Il n'est pas recommandé de réaliser systématiquement une scintigraphie osseuse en cas de cancer de la prostate localisé à risque faible.	Isotopen botskans zijn niet routinematig aanbevolen bij mannen met laag-risico gelokaliseerd prostaatCA.
4. Petscan	Il n'est pas recommandé de réaliser un examen par émission de positron (PET) en cas de cancer de la prostate localisé à risque faible.	Een PET-scan is niet aanbevolen bij mannen met laag-risico gelokaliseerd prostaatCA.
5.Surveilact	La surveillance active comprenant la possibilité d'un traitement différé est une stratégie valide dans les cancers localisés à risque faible.	Actieve opvolging met uitstel van behandeling is een valide strategie voor gelokaliseerd laag-risico prostaatCA.
6. IRM	L'IRM et les marqueurs tumoraux (PCA3 et PSA PHI) sont des stratégies efficaces dans la surveillance active des cancers indolents.	IRM and PCA3-PSA PHI vormen een valide strategie voor actieve opvolging van het “indolent” (slapend) prostaatCA.
7. Prostatectomy	La prostatectomie radicale et la radiothérapie externe ou interne (brachythérapie) aboutissent aux mêmes résultats d'un point de vue oncologique en cas de cancer localisé à risque faible ou intermédiaire.	Radicale prostatectomie, LDR of HDR brachytherapie en EBRT geven gelijkaardige oncologische resultaten bij patiënten met een laag- of intermediair risico, gelokaliseerd prostaatCA.
8. Prostatectomy_risk	La prostatectomie radicale et la radiothérapie externe ou interne (brachythérapie) entraînent les mêmes risques de problèmes fonctionnels (dysfonction érectile et incontinence) en cas de cancer localisé à risque faible ou intermédiaire.	Radicale prostatectomie, LDR of HDR bracytherapie en EBRT geven gelijkaardige functionele resultaten (erectie, continence) bij patiënten met een laag- of intermediair risico, gelokaliseerd prostaatCA.
9. Prostatec_oncoc	La prostatectomie aboutit aux mêmes résultats d'un point de vue oncologique, quelque soit la technique utilisée (prostatectomie à ciel ouvert, radicale ou assistée par robot) en cas de cancer localisé à risque faible ou intermédiaire.	Open, laparoscopische of robot-geassisteerde radicale prostatectomie geven gelijkaardige oncologische resultaten bij patiënten met een laag- of intermediair risico, gelokaliseerd prostaatCA.
10. Prostatec-alltypes	La prostatectomie entraine les mêmes risques de problèmes	Open, laparoscopische of robot-geassisteerde radicale



	fonctionnels (dysfonction érectile et incontinence) en cas de cancer localisé à risque faible ou intermédiaire, quelque soit la technique utilisée (prostatectomie à ciel ouvert, radicale ou assistée par robot).	prostatectomie geven gelijkaardige functionele resultaten (erectie, continentie) bij patiënten met een laag- of intermediair risico, gelokaliseerd prostaatCA.
11. HIFU	Le traitement par HIFU n'est pas recommandé en dehors des études cliniques contrôlées en cas de cancer localisé.	HIFU is niet aanbevolen bij patiënten met gelokaliseerd prostaatCA buiten de context van een gecontroleerde klinische studie.
12. Hormonothérapie	Une hormono-thérapie (néo)adjuvante est indiquée en combinaison à la radiothérapie externe ou interne (brachythérapie).	(Neo-)adjuvante hormonale therapie is aangewezen in combinatie met EBRT en brachytherapie.
13. Hormono_Highriskcancer	Une thérapie hormonale (néo)adjuvante est indiquée en combinaison à la radiothérapie externe ou interne (brachythérapie) en cas de cancer localisé à haut risque.	(Neo-)adjuvante hormonale therapie is aangewezen in combinatie met EBRT en brachytherapie bij patiënten met hoog-risico gelokaliseerd prostaatCA.
14. Hormono_prostatec	Une thérapie hormonale adjuvante n'est pas indiquée en combinaison à la prostatectomie radicale totale en cas de cancer localisé à haut risque.	Adjuvante hormonale therapie is niet aangewezen in combinatie met radicale prostatectomie bij patiënten met hoog-risico gelokaliseerd prostaatCA.
15. Hormono_étatpatients	Il est envisageable de traiter par hormonothérapie les patients ayant une tumeur localisée qui ne sont pas en état de subir un traitement local.	Patiënten met gelokaliseerd prostaatCA die niet geschikt zijn voor lokale therapie moeten behandeld worden met hormonale therapie.
16. Hormono_étatpatients_avancés	Il est envisageable de traiter par hormonothérapie les patients ayant une tumeur localement avancée qui ne sont pas en état de subir un traitement local.	Patiënten met gelokaliseerd gevorderd prostaatCA die niet geschikt zijn voor radicale therapie, moeten behandeld worden met hormonale therapie.
17. Discuss_options	Il convient que les soignants discutent des différentes options de traitement avec les patients.	Zorgverstrekkers moeten alle relevante behandelingsstrategieën overleggen met mannen met prostaatCA.
18. Info_effets	Il convient que les soignants informent les patients des effets du cancer de la prostate et des conséquences potentielles des différents traitements en matière de continence et de sexualité.	Zorgverstrekkers moeten adequate informatie verstrekken aan mannen met prostaatCA aangaande de effecten van prostaatCA en van de behandelingsopties op de seksuele functie en op de continentie.
19. Discuss_uro§radio	Vu qu'il existe plusieurs options de traitement et qu'elles peuvent entraîner des effets secondaires sérieux, il convient que les patients qui sont candidats à un traitement radical aient la possibilité de discuter des traitements avec un urologue et un radiothérapeute.	Gegeven de brede waaier van behandelingsmodaliteiten en hun ernstige nevenwerkingen, moeten mannen met prostaatCA die kandidaten voor radicale therapie zijn de kans krijgen om overleg te plegen met een uroloog en een radiotherapeut.
20. Hormono_récidive	En cas de récurrence (PSA) après prostatectomie radicale, un traitement hormonal (de « sauvetage ») est indiqué.	In geval van PSA recidief na radicale prostatectomie is « salvage » hormonale therapie aangewezen.



Appendix 2.2. Websurvey “prostate” results

1. Groupe-cible	Fréquences	Pourcentages
Urologue	54	60,0%
Radiothérapeute	13	14,4%
Généraliste	10	11,1%
Infirmier(ère) en urologie	3	3,3%
Patient	10	11,1%
Total/ répondants	90	

Interrogés : 90 / Répondants : 90 / Réponses : 90. Pourcentages calculés sur la base des répondants

2. CTscanpelvis	Fréquences	Pourcentages
D'accord	48	53,9%
Pas d'accord	41	46,1%
Total	89	100,0%

3. Scintigraphie	Fréquences	Pourcentages
D'accord	79	88,8%
Pas d'accord	10	11,2%
Total	89	100,0%



4. Petscan	Fréquences	Pourcentages
D'accord	88	98,9%
Pas d'accord	1	1,1%
Total	89	100,0%

5. Surveilact	Fréquences	Pourcentages
D'accord	83	93,3%
Pas d'accord	6	6,7%
Total	89	100,0%

6. IRM	Fréquences	Pourcentages
D'accord	38	44,7%
Pas d'accord	47	55,3%
Total	85	100,0%

7. Prostatectomy	Fréquences	Pourcentages
D'accord	58	65,9%
Pas d'accord	30	34,1%
Total	88	100,0%



8. Prostatectomy_risk	Fréquences	Pourcentages
D'accord	20	23,0%
Pas d'accord	67	77,0%
Total	87	100,0%

9. Prostatectomy_onco	F Fréquences	Pourcentages
D'accord	69	79,3%
Pas d'accord	18	20,7%
Total	87	100,0%

10. Prostatectomy_alltypes	Fréquences	Pourcentages
D'accord	45	51,1%
Pas d'accord	43	48,9%
Total	88	100,0%

11. HIFU	Fréquences	Pourcentages
D'accord	83	94,3%
Pas d'accord	5	5,7%
Total	88	100,0%



12. Hormonothérapie	Fréquences	Pourcentages
D'accord	27	31,4%
Pas d'accord	59	68,6%
Total	86	100,0%

13. Hormono_highriskcancer	Fréquences	Pourcentages
D'accord	79	89,8%
Pas d'accord	9	10,2%
Total	88	100,0%

14. Hormono_prostatectomy	Fréquences	Pourcentages
D'accord	57	67,1%
Pas d'accord	28	32,9%
Total	85	100,0%

15. Hormono_étatpatients	Fréquences	Pourcentages
D'accord	36	41,4%
Pas d'accord	51	58,6%
Total	87	100,0%



16. Hormono_étatpatients_avancé	Fréquences	Pourcentages
D'accord	75	85,2%
Ppas d'accord	13	14,8%
Total	88	100,0%

17. Discuss_options	Frequency	Pourcentages
D'accord	88	98,9%
Pas d'accord	1	1,1%
Total	89	100,0%

18. Info_effets	Fréquences	Pourcentages
D'accord	88	100,0%
Total	88	100,0%

19. Discuss_uro&radio	Fréquences	Pourcentages
D'accord	69	78,4%
Pas d'accord	19	21,6%
Total	88	100,0%



20. Hormono_récidive	F Fréquences	Pourcentages
D'accord	36	41,9%
Pas d'accord	50	58,1%
Total	86	100,0%

22. Language	Fréquences	Pourcentages
FR	37	41,1%
NL	53	58,9%
Total	90	100,0%



APPENDIX 3. LITERATURE SEARCH

Appendix 3.1. Clinical practice guidelines

Appendix 3.1.1. Sources

A broad search of electronic databases (Medline, EMBASE), specific guideline websites and websites of oncologic organisations (Table 7) was conducted in February 2011. Only guidelines published in Dutch, English, French or German and after 01/01/2005 were selected.

Table 7–Searched guideline websites and websites of oncologic organisations

0	Alberta Heritage Foundation For Medical Research (AHFMR)	http://www.ahfmr.ab.ca/
0	American Society of Clinical Oncology (ASCO)	http://www.asco.org/
0	American College of Surgeons (ACS)	http://www.facs.org/cancer/coc/
5	CMA Infobase	http://mdm.ca/cpgsnew/cpgs/index.asp
15	Guidelines International Network (GIN)	http://www.g-i-n.net/
1	National Comprehensive Cancer Network (NCCN)	http://www.nccn.org/
1 new, 5 ACR	National Guideline Clearinghouse	http://www.guideline.gov/
0	National Cancer Institute	http://www.cancer.gov/
1 duplicate	Haute Autorité de Santé (HAS)	http://bfes.has-sante.fr/HTML/indexBFES_HAS.html
0	BC Cancer Agency	http://www.bccancer.bc.ca/default.htm
10 on screening	Institute for Clinical Systems Improvement (ICSI)	http://www.icsi.org/index.asp
1 (all)	National Health and Medical Research Council (NHMRC)	http://www.nhmrc.gov.au/
0	Scottish Intercollegiate Guidelines Network (SIGN)	http://www.sign.ac.uk/
2 duplicate	New Zealand Guidelines Group (NZGG)	http://www.nzgg.org.nz/
1 HAS + update (all)	Fédération Nationale des Centres de Lutte Contre le Cancer (FNCLCC)	http://www.fnclcc.fr/sor/structure/index-sorspecialistes.html
5 duplicate	National Institute for Health and Clinical Excellence (NICE)	http://www.nice.org.uk/



Appendix 3.1.2. Search terms for Medline (Ovid)

For Medline the following MeSH or non MESH terms were used in combination: "prostatic neoplasms"[MeSH Terms] OR prostate cancer[Text Word]. For EMBASE the following Emtree terms were used in combination: "prostate cancer, watchful waiting". These MeSH and Emtree terms were combined with a standardised search strategy to identify CPGs (Table 8).

Table 8 – Standardised search strategy for CPGs

Database	Search strategy
Medline	guideline [pt] OR practice guideline [pt] OR recommendation* [ti] OR standard* [ti] OR guideline* [ti]
EMBASE	'practice guideline'/exp

Exclusion criteria : advanced stage :

Medline (Ovid) search :

prostatic neoplasms.mp. or Prostatic Neoplasms/ (73381)

2 prostate cancer.mp. (47158)

3 Guideline/ or Practice Guideline/ (20402)

4 "recommendation*".m_titl. (17706)

5 "standard*".m_titl. (52575)

6 "guideline*".m_titl. (36194)

7 1 or 2 (79090)

8 3 or 4 or 5 or 6 (113470)

9 7 and 8 (632)

10 limit 9 to (humans and yr="2005 -Current" and (dutch or english or french or german)) (256)

Appendix 3.1.3. Results

After exclusion of duplicates, 20 guidelines were found on specific website. Two hundred fifty six publications were retrieved on Pubmed (Ovid). Based on title and abstract and after exclusion of duplicates, 17 publications issued from Ovid were selected. Finally, we found 37 guidelines. Some guidelines are focused on specific questions. Therefore, they are presented by topics on Table 9. First column shows guidelines what take all the prostate cancer process into account. Second column shows guidelines focused on prostate cancer diagnosis. Third column shows guidelines focused on all prostate cancer treatment. Column five shows guidelines focused on prostate cancer follow-up, column six on prostate cancer by men > 70y and column seven on news technologies.



Table 9 – All guidelines found by topics (first selection)

All	Diagnosis			Treatment				T3	Follow-up	Senior 70 y	HIFU/cryo
	Biopsy	PET/FDG	Bone scan	Radical	Brachy	IMRT	Combined				
NICE 2008	NICE 2010	Bourguet 2007	Briganty		ACR	Maceira Rozas 2006	NZGG	Casas		Droz JP	
NovaScotia 2006	Bertaccini 2007			CCO	NICE Low	CCO	Sidhom 2008		McIntosh	Mongiat-Arthus	CCO
Saska 2008	CMA 2009				NICE High (174)		Hoffman		Richaud 2005		NICE
EAU 2010 (Heidenrich)					IQWIG				Roach		NICE
AUA (20)					Kovacks 2005						
VIKC, De Reijke 2008											
ESMO, Horwich 2008											
Kamidono 2008											
NCCN 2010											
S3, Wenz											

We excluded for this part 26 guidelines outside of the scope. We performed first a rapid appraisal of 11 guidelines quality based on questions 7,8 and 10 included in AGREE tool. Those questions are focused on selecting the evidence .



Table 4 – Rapid appraisal based on Questions 7, 8, 10 of AGREE (version May 2009 and scale 1 to 7)

Reference	Q 7 : Systematic methods were used to search evidence	Q 8 : The criteria for selecting the evidence were clearly described	Q 10 : The methods used for formulating recommendations were clearly described	Conclusion
NICE 2008 Made by NCC-C Lit search: 1 June 2007	7	7	7	21/21, included
NovaScotia 2006	1	1	1	Methodology not described excluded
Saskatchewan Cancer Agency Update 2008	1	1	1	Methodology not described, decisions based on consensus excluded
AUA 2007 Lit search: April 2004	7	6	6	19/21, included
HAS, (SOR)update 2008	6	1	1	Excluded (after expert discussion)
EAU 2010 Lit search: update janv 2010	6	4	5	15/21: tentatively included
VIKC, De Reijke 2008	5	6	6	19/21, included
ESMO	1	1	1	Methodology not described excluded
Kamidono	5	6	3	14/21: excluded
NCCN (2010)	1	1	1	Excluded
Aragon (2011)	6	4	4	14/21: tentatively included



We considered a good evidence selecting process as a basic requirement and decided not to follow the appraisal if no good description of this process was available. Then, we performed the appraisal of the 5 (tentatively) included guidelines with the entire AGREE tool (see point 4.1).

Appendix 3.2. Systematic reviews and primary studies for question 1

- P: localized prostate cancer
- I: active surveillance or watchful waiting
- C: other treatment
- O : mortality, morbidity, quality of life

Appendix 3.2.1. Medline

Keywords

- MeSH : Prostatic Neoplasms :Tumours or cancer of the PROSTATE.
- Prostatic Neoplasms/th : therapy of Tumours or cancer of the PROSTATE
- PROSTATECTOMY : Complete or partial surgical removal of the prostate. Three primary approaches are commonly employed: suprapubic - removal through an incision above the pubis and through the urinary bladder; retropubic - as for suprapubic but without entering the urinary bladder; and transurethral (TRANSURETHRAL RESECTION OF PROSTATE).
- Text: *Watchful Waiting

Project number	2011-01-GCP
Project name	Prostate cancer
Search question(s)	Role of active surveillance/watchful waiting in the management of localised prostate cancer (low, intermediate and high risk)?
Structured search question(s) (PICO, and related keywords SPICE, ECLIPSE, ..)	

P (patient)	Patient with localised prostate cancer	“Prostatic neoplasms” or local\$ prostate cancer
I (Intervention)	active surveillance/watchful waiting	*Watchful Waiting or active surveillance
C (comparison)	Surgery or other treatment	Prostatectomy/
O (outcome)	Mortality, morbidity, quality of life	

Date	07/12/2011
Database	Ovid MEDLINE(R) without Revisions 1996 to November Week 3 2011
Search Strategy	Prostatic Neoplasms/ (54530) 2 local\$ prostate cancer.mp. (3900) 3 1 or 2 (54655) 4 watchful waiting.mp. or *Watchful Waiting/ (1289) 5 androgen antagonists.mp. or Androgen Antagonists/ (5198) 6 brachytherapy.mp. or Brachytherapy/ (10409) 7 neoadjuvant therapy.mp. or Neoadjuvant Therapy/ (8440) 8 prostatectomy.mp. or Prostatectomy/ (14223) 9 radiotherapy.mp. or Radiotherapy, Intensity-Modulated/ or Radiotherapy/ or Radiotherapy, Conformal/ (84353) 10 5 or 6 or 7 or 8 or 9 (110055) 11 exp Controlled Clinical Trial/ (37584) 12 exp Randomized Controlled Trials as Topic/



(67539)
 13 exp Random Allocation/ (37598)
 14 exp Double-Blind Method/ (67080)
 15 exp Single-Blind Method/ (12779)
 16 exp Clinical Trial/ (446179)
 17 (clin\$ adj25 trial\$).tw. (152122)
 18 ((single or double or triple or treble) adj25
 (blind\$ or mask\$)).tw. (67282)
 19 placebo\$.tw. (85758)
 20 random\$.tw. (402126)
 21 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or
 19 or 20 (822281)
 22 (watchful waiting\$ or observation\$ or expectant
 management\$ or careful monitoring\$ or
 surveillance\$ or delay\$ or deferred treatment\$ or
 no initial treatment\$).tw. (477802)
 23 (watch and wait\$).tw. (265)
 24 22 or 23 (478011)
 25 4 or 24 (478117)
 26 3 and 10 and 21 and 25 (455)

Note 18 (5 on quality of life) remained before critical appraisal

Appendix 3.2.2. Cochrane Library

Date	From 2007 until 07/01/2011
Database	CDSR, DARE, CCRCT
Search Strategy (MESH term)	Prostatic Neoplasms/th
Note	CDSR (6), DARE(5), CCRCT (60), total : 71

Appendix 3.2.3. Embase

Entree terms :

- prostate cancer, castration resistant prostate cancer, prostate adenocarcinoma, prostate carcinoma, prostatic intraepithelial neoplasia, (This term was added to Emtree in 1974),Synonyms : cancer, prostate; prostate gland cancer; prostatic cancer
- “watchful waiting” . This term was added to Emtree in 2007
- Active surveillance: no emtree term, free text
- Prostatectomy: This term was added to Emtree in 1974, Synonyms:prostate adenectomy; prostate resection; prostatic adenectomy; radical prostatectomy; total prostatectomy. **Dorland's dictionary:** prostatectomy = surgical removal of the prostate or of a part of it.Radical prostatectomy = removal of the prostate with its capsule, seminal vesicles, ductus deferens, some pelvic fasciae, and sometimes pelvic lymph nodes; performed via either the retropubic or the perineal route.




- Radiotherapy . This term was added to Emtree in 1974. Synonyms : bioradiant therapy; bucky irradiation; bucky radiation; bucky radiotherapy; bucky ray; bucky ray radiation; bucky therapy; fractionated radiotherapy; hemibody irradiation; hypophysectomy, radiation; hypophysis irradiation; hypophysis radiation; irradiation therapy; irradiation treatment; irradiation, hypophysis; lymphatic irradiation; pituitary irradiation; radiation beam centration; radiation repair; radiation therapy; radiation treatment; radio therapy; radio treatment; radiohypophysectomy; radiology, therapeutic; radiotreatment; roentgen irradiation, therapeutic; roentgen therapy; roentgen treatment; therapeutic radiology; therapy, irradiation; therapy, radiation; therapy, roentgen; treatment, irradiation; treatment, radiation; treatment, roentgen; x-ray therapy; x radiotherapy; x ray therapy; x ray treatment. Emtree Scope Note radiotherapy = Used as a disease subheading for the treatment of a disease using radiotherapy
- (Neo)adjuvant therapy. This term was added to Emtree in 1974. Synonyms : adjuvant effect; adjuvant treatment; neoadjuvant therapy; radiotherapy, adjuvant. **Dorland's dictionary** : neoadjuvant therapy = in combined modality therapy for cancer, initial use of one modality, such as chemotherapy or radiotherapy, to decrease the tumour burden prior to treatment by another modality, usually surgery. Called also preoperative t. and presurgical t.
- adjuvant therapy = the use of chemotherapy or radiotherapy in addition to surgical resection in the treatment of cancer.
- Androgen antagonist . This term was added to Emtree in 1974. Synonyms : androgen antagonist; androgen antagonists; anti androgen; antiandrogen agent; antiandrogenic agent; antiandrogenic drug; nonsteroidal anti-androgen; nonsteroidal anti-androgens; nonsteroidal anti androgen; nonsteroidal anti androgens; nonsteroidal antiandrogen; nonsteroidal antiandrogens. **Dorland's dictionary** : antiandrogen = any substance capable of inhibiting the biological effects of androgens.

Date	From 2007 until 01/2011
Database	Embase
Search Strategy (Emtree terms, see below)	'prostate cancer'/exp AND 'watchful waiting'/exp AND (watchful AND waiting OR observation OR expectant AND management OR careful AND monitoring OR surveillance OR delay OR (deferred AND treatment OR no AND initial AND treatment)) AND ('prostatectomy'/exp OR 'radiotherapy'/exp OR 'antiandrogen'/exp OR 'adjuvant therapy'/exp) AND [embase]/lim AND [2007-2012]/pyAND 'clinical trial'/de OR 'controlled clinical trial'/de OR 'controlled study'/de OR 'double blind procedure'/de OR 'human'/de OR 'prospective study'/de OR 'randomized controlled trial'/de
Note	55 (16 duplicates with Medline)



Appendix 3.2.4. Results



The quality of the retrieved SR, M-A and RCT was assessed using the checklists of the Dutch Cochrane Centre (www.cochrane.nl). All critical appraisals were done by a single KCE expert.

From those 22 relevant studies, 9 were included after critical appraisal (see point 4.1 and 4.2) and 13 not (see point 4.3)

Appendix 3.3. Randomised control trials for question 3

P: localized prostate cancer
 I: protocol X for active surveillance
 C: protocol Y for active surveillance
 O: mortality, morbidity, quality of life

Appendix 3.3.1. Medline

Project number	2011-01-GCP
Project name	Prostate cancer /active surveillance strategies
Search question(s)	Did different active surveillance strategies affect the outcomes? (question 4.3.1)
Structured search question(s) (PICO, SPICE, ECLIPSE, ..)	and related keywords
P (patient)	Low risk prostate cancer Prostatic Neoplasms
I (Intervention)	Protocol1 Active surveillance
C (comparison)	Protocol2
O (outcome)	mortality
Date	09/08/2012
Database	Database: Ovid MEDLINE(R) < 2008 to August Week 1 2012>
Search Strategy	1 exp Prostatic Neoplasms/ (20765) 2 Prostatic Intraepithelial Neoplasia/ (279) 3 (prostat\$ adj3 (cancer\$ or carcinoma\$ or adeno\$ or malignan\$ or tum?\$ or neoplas\$ or intraepithelial\$)).tw. (21304) 4 PIN.tw. (1811)



-
- 5 1 or 2 or 3 or 4 (26441)
 - 6 watchful wait\$.tw. (370)
 - 7 (watch\$ adj2 wait\$).tw. (494)
 - 8 watchful observation.tw. (5)
 - 9 watchful surveillance.tw. (2)
 - 10 watchful monitoring.tw. (6)
 - 11 active surveillance.tw. (1136)
 - 12 active monitoring.tw. (63)
 - 13 expectant manag\$.tw. (350)
 - 14 expectant monitoring.tw. (13)
 - 15 expectant surveillance.tw. (1)
 - 16 deferred treatment\$.tw. (23)
 - 17 deffered therap\$.tw. (0)
 - 18 delayed treatment\$.tw. (440)
 - 19 delayed therap\$.tw. (60)
 - 20 conservative monitoring.tw. (2)
 - 21 conservative surveillance.tw. (1)
 - 22 6 or 7 or 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 (2509)
 - 23 5 and 22 (508)
 - 24 limit 23 to yr="2011 -Current" 38
-

Appendix 3.3.2. Cochrane

Date	From 2008 until 09/08/2012
-------------	----------------------------

Database	CDSR, CCRCT
-----------------	-------------

Search Strategy (MESH term)	Prostatic Neoplasms/th
------------------------------------	------------------------

Result	After exclusion of articles non relevant for this question : no
---------------	---

Appendix 3.3.3. Embase

Date	From 2008 until 10/08/2012
-------------	----------------------------

Database	Embase
-----------------	--------

Search Strategy (Emtree terms, see below)	'prostate cancer'/exp/mj OR 'prostate cancer' AND active AND surveillance AND [controlled clinical trial]/lim AND [english]/lim AND [2011-2013]/py
--	--

Result	7
---------------	---

Forty four potentially relevant citations were identified.

After reviewing the 11 publications selected, no studies comparing different active surveillance strategies was found.



APPENDIX 4. QUALITY APPRAISAL

Appendix 4.1. Guidelines

Table 5 – Guidelines appraisal following AGREE tool

	NICE	AUA	EAU	VIKC-CBO	ARAGON
Domain 1. Scope and Purpose					
1. overall objective	6,5	7,0	6,0	7,0	6,5
2. health question(s)	6,5	6,5	6,0	7,0	7,0
3. population	7,0	7,0	6,0	6,5	7,0
Domain score	94,4	97,2	83,3	97,2	100,0
Domain 2. Stakeholder Involvement					
4. all relevant professional groups	6,5	6,5	5,0	6,0	4,5
5. target population views and preferences	6,0	1,0	1,0	3,5	1,0
6. target users	7,0	6,0	6,5	7,0	4,0
Domain score	91,7	58,3	52,8	75,0	50,0
Domain 3. Rigour of Development					
7. systematic search	7,0	7,0	6,0	4,5	6,0
8. selection criteria	6,0	6,5	3,5	5,0	4,0
9. strengths and limitations of evidence	5,0	6,5	5,5	4,0	4,5

10. formulation of recommendations	6,0	6,0	5,0	6,0	4,0
11. benefits, side effects and risks	6,0	6,0	5,0	5,5	2,5
12. explicit link	6,5	6,0	7,0	6,5	4,0
13. external review	5,5	5,0	1,0	5,0	5,0
14. update procedure	6,5	5,5	5,5	6,0	6,5
Domain score	84,4	84,4	56,3	67,7	70,8
Domain 4. Clarity of Presentation					
15. specific and unambiguous	6,5	6,5	5,5	4,0	7,0
16. different options for management	7,0	7,0	6,0	6,0	4,0
17. key recommendations	7,0	7,0	5,5	7,0	6,5
Domain score	97,2	97,2	77,8	77,8	100,0
Domain 5. Applicability					
18. facilitators and barriers	5,0	1,0	1,0	7,0	1,5
19. advice and tools	5,0	3,0	1,0	6,5	6,0
20. possible resource implications	1,0	1,0	1,0	1,0	1,0
21. monitoring and/or auditing criteria	4,0	1,0	1,0	6,0	1,0
Domain score	45,8	8,3	0,0	68,8	25,0
Domain 6. Editorial Independence					



22. editorially independent	6,5	6,5	6,5	6,5	6,5
23. conflicts of interest	7,0	6,5	6,5	6,5	7,0
Domain score	95,8	91,7	91,7	91,7	100,0
Number of items scoring ≥ 5	21	18	16	18	11
Number of domains scoring > 60%	5	4	3	6	4

Finally, we selected three guidelines⁸⁻¹⁰ of high quality and two^{11, 26} of moderate quality according to AGREE score.

Appendix 4.2. Systematic Reviews

Items	AHRQ	Farmaka	Hegarty	Wilt-SR
Search date	August 2011	August 2010	July 2010	September 2007
Inter-vention	Curative Treatment	Radical Prostatectomy (RP)	Curative treatment	Any prostate cancer treatment
Control	AS	WW/AS	WW	Any prostate cancer treatment
1	Yes	Yes	Yes	Yes
2	No	: Yes/No	Yes	Yes

	restricted			
3	?	?	Yes	Yes
4	Yes/No	Yes/No	Yes	Yes
5	Yes	Yes	Yes	Yes
6	Yes	Yes	Yes	Yes
7	NA	NA	NA	NA
8	NA	NA	NA	NA
9	Yes	Yes	Yes	Yes
Quality	Moderate	Moderate	High	High

- Legend of items 1 to 9 of the quality appraisal:*
1. Is de vraagstelling adequaat geformuleerd?
 2. Is de zoekactie adequaat uitgevoerd?
 3. Is de selectieprocedure van artikelen adequaat uitgevoerd?
 4. Is de kwaliteitsbeoordeling adequaat uitgevoerd?
 5. Is adequaat beschreven hoe data-extractie heeft plaatsgevonden?
 6. Zijn de belangrijkste kenmerken van de oorspronkelijke onderzoeken beschreven?
 7. Is adequaat omgegaan met klinische en statistische heterogeniteit van de onderzoeken ?
 8. Is statistische pooling op een correcte manier uitgevoerd ?
 9. Zijn de resultaten van de systematische review valide en toepasbaar?



Appendix 4.3. Randomised Controlled Trials

Items	Bill-Axelsson	Iversen	Fransson	Johansson	Wilt-PIVOT
Intervention	RP	RP + oral placebo	Radiotherapy	RP	RP
Control	WW	oral placebo alone	WW	WW	Observation
1	Yes	Yes	?	Yes	Yes
2	Yes	?	?	Yes	?
3	No	No	?	No	No (not possible)
4	No	No	?	No	No (not possible)
5	Yes	?	?	Yes	?
6	Yes	No	Yes	No	Yes
6 bis		No		?	
7	Yes	No	?	?	Yes
7 bis		No		?	
8	Yes	Yes	Yes	Yes	Yes
9	Yes	Yes		Yes	Yes
10	Non car le cancer de la prostate est détecté par le dosage de PSA actuellement, ce qui n'est pas le cas dans cette RCT.	Douteux car de nombreuses questions méthodologiques entachent les résultats : pouvoir statistique, exclusion de l'analyse de 22% des hommes de l'échantillon initial, différence d'âge entre les 2 groupes, diagnostic sans bone scan.	Douteux. Trop peu d'information dans cet article pour répondre à cette question.	Non, les résultats sont peu applicables au vu du mode de diagnostic (non basé sur le dosage de PSA) et de la chirurgie peu conservative pratiquée.	Oui, beaucoup plus proche de la situation nationale car $\frac{3}{4}$ de la population a été diagnostiquée à la suite d'un PSA élevé.
Quality	High	Poor	poor	High	high
	<p>1. L'attribution de l'intervention aux patients a-t-elle été réalisée de manière aléatoire (randomisation)?</p> <p>2. Celui qui inclut les patients ignore l'ordre de succession aléatoire. Est-ce le cas?</p> <p>3. Le traitement était-il étudié en aveugle pour le patient?</p> <p>4. Le traitement était-il réalisé en aveugle pour les soignants?</p>	<p>5. Le traitement était-il réalisé en aveugle pour ceux qui évaluent les résultats ?</p> <p>6. Les groupes étaient-ils comparables au début de l'étude ?</p> <p>6 bis. Si non, cela a-t-il été corrigé lors de l'analyse des résultats ?</p> <p>7. Parmi les patients inclus, une proportion suffisante a-t-elle disposé d'un suivi complet ?</p>	<p>7 bis. Si non : une perte sélective de patients lors du suivi est-elle exclue?</p> <p>8. Tous les patients inclus ont-ils été analysés dans les groupes dans lesquels ils ont été randomisés ?</p> <p>9. En dehors de l'intervention, les groupes sont-ils traités de façon semblable ?</p> <p>10. Le résultat obtenu peut-il être appliqué à la situation nationale ?</p>		



APPENDIX 5. DATA EXTRACTION TABLES

Appendix 5.1. Systematic reviews

All systematic reviews included the same two controls trials: the SPCG-4 trial (good quality) (see Bill-Axelsson point 5.2.1) and the VACURG (poor quality) (see Iversen point 5.2.1).

I Study ID	II Method	III Patient characteristics	IV Intervention(s)	V Results primary outcome	VI Results secondary and other outcome(s)	VII Critical appraisal of review quality
<p>AHRQ. “An Evidence Review of Active Surveillance in Men With Localized Prostate Cancer”</p>	<p>Systematic review to summarize the existing literature regarding the role of AS in the management of early-stage, low-risk prostate cancer</p> <p>(1 subquestion is comparative effectiveness of AS vs active treatments).</p> <p>Funding : National Institutes of Health (NIH) Office of Medical Applications of Research (OMAR)</p> <p>Search date: August 2011</p> <p>Database: Medline (OVID), CDSR, specific database (epidemiology, economic) et experts</p> <p>Study designs : RCT & non randomize comparative studies of treatments, multivariable association studies, & studies of temporal trends in prostate cancer natural history. Only published, peer-reviewed, English-language articles were selected based on predetermined eligibility</p>	<p>Men with clinically localized prostate cancer (T1-T2), without known lymph nodes (N0-X) or metastases (M0-X)</p>	<p>Radical prostatectomy (RP) or external beam radiation therapy (RT) or brachytherapy (alone or combined), with or without ADT.</p> <p>Active surveillance (AS) = In fact, there was no standardized definition of active surveillance. Sixteen cohorts used different monitoring protocols, all with different combinations of periodic digital rectal examination, PSA testing, rebiopsy, and/or imaging findings. Predictors that a patient received no initial active treatment generally included older age, presence of comorbidities, lower Gleason score, lower tumour stage, lower diagnostic PSA, and lower disease progression risk group.</p>	<p>No trial provided results comparing men with localized disease on active surveillance with surgery or radiation therapy.</p> <p>Therefore, there is insufficient evidence for the comparative short- and long-term outcomes of AS versus immediate definitive treatment for localized prostate cancer.</p> <p>For other observational management strategies (largely resembling WW), in addition to previously published systematic reviews and evidence reports, 2 updates from multicenter RCTs and 16 cohort studies (3 prospective and 13 retrospective) were selected.</p> <p>- lower all-cause or prostate cancer-specific mortality rates with RP vs WW; SPCG4 (see Bill-Axelsson point 4.2.1) trial found significantly lower incidences of all-cause deaths (24 vs. 30 percent), disease-specific deaths (10 vs. 15 percent),</p>	<p>- less risk of urethral stricture with observational strategy vs RP;</p> <p>- higher short and long term cost for RP or RT vs WW.</p>	<p>Moderate</p> <p>The literature search focused on Medline & CDSR only and it's not clear if the quality appraisal of the primary studies was taken into account.</p> <p>The majority of evidence for this Key Question came from retrospective analyses of observational studies. Confounding by indication is likely in these studies, due to the differences in patient characteristics and risk profile between patients treated with observational strategies and those who received active treatment.</p> <p>No pooling was done</p>



	criteria No trial found for the sub question about the effectiveness of AS but 2 updates RCT, 16 cohort studies for other observational management strategies (largely resembling WW), 4 reports and 2 cost modelin					
Farmaka Le traitement des pathologies prostatiques bénignes et malignes 2011	<p>Systematic review to:</p> <ul style="list-style-type: none"> - prepare a consensus conference about the "traitement effi cient des pathologies bénignes et malignes de la prostate" - a subquestion of this search was : "Dans quels cas (criteres precis) une attitude de surveillance active peut-elle etre justifiee pour un cancer de la prostate ?" <p>Funding : National Sickness Fund</p> <p>Search date: August 2010</p> <p>Database: MEDLINE, Cochrane Library & Dare.</p> <p>Study designs : 1 RCT & 7 Observational study. Three ongoing studies were cited .</p>	Men with prostate cancer (no other precision)	<p>Radical prostatectomy</p> <p>WW= to avoid treatment; palliative intent</p> <p>AS= to individualize treatment; curative intent</p>	<p>There are currently no published RCT on active surveillance (only on watchfull waiting).</p> <p>In the RCT on WW (SPCG-4, sse Bill-Axelsson point 4.2.1)), WW is associated with a slight specific mortality rate and a reduced local and metastatic progression.</p> <p>In the 7 cohort studies (with a maximum follow-up of 7 years), AS in men with early stage PCa is associated with slight specific (> 3% at max 13 years of follow-up) mortality rates.</p> <p>The exact place of AS, the type of patients who get the greatest benefit of this approach and the monitoring protocol are to be defined.</p>		<p>Moderate</p> <p>ADAPTE methodology</p> <p>No pooling was done</p>
Hegarty J. & al: "Radical prostatectomy versus watchful waiting for prostate"	<p>Systematic review to:</p> <ul style="list-style-type: none"> - compare the beneficial and harmful effects of RP versus WW for the treatment of clinically localised prostate cancer - test the null hypothesis of no difference in terms of the primary, secondary 	Men with clinically localized prostate cancer = confirmed prostate cancer (as verified by cytological	Radical prostatectomy (RP) = removal of the entire prostate gland and some surrounding tissue performed by any method (e.g. retropubic, perineal, laser, robotic or laparoscopic) with or without nerve sparing	<p>All cause mortality after 23 years (VACURG):</p> <p>10.6 years for RP vs 8 years for WW (p>0.05) (but poor methodological quality of the trial)</p> <p>All cause mortality after 15 years (VACURG):</p> <p>Hazard Ratio (HR) = 0.9 (0.56</p>	<p>Distant metastases after 12 years (SPCG-4) :</p> <p>19.3% vs 26%</p> <p>RD = -6.7% (-13.2 to -0.2);</p> <p>RR = 0.65 (0.47 to 0.88);</p> <p>p=0.006</p> <p>Incidence of local recurrence and/or progression after 12</p>	<p>High</p> <p>None RCT enrolled men with primarily PSA-detected disease (only 5.2% in SPCG-4)</p> <p>Not applicable to men with significant co-morbidities (enrollment of men < 75 years with a LE of 10 years in SPCG-4)</p>



<p>cancer (Review)"</p>	<p>and tertiary outcomes between RP and WW Funding : Health Research Board, Ireland Search date: 30 July 2010 Database: MEDLINE, EMBASE, The Cochrane Library, ISI Science Citation Index, DARE, LILACS. Study designs : RCT or quasi-RT</p>	<p>or histological examination) which is believed to be still confined to the prostate gland (T0 or T1 or T2).</p>	<p>procedures, with or without sampling of the pelvic lymph nodes. Watchful waiting (WW) = any conservative approach whereby a decision is made to provide no initial treatment and to monitor the patient. Palliative treatment can offer if evidence of disease progression.</p>	<p>to 1.43). All cause mortality after 12 years (SPCG-4): 32.7% vs 39.8% RD = -7.1% (-14.7 to 0.5); RR = 0.82 (0.65 to 1.03); p=0.09 Prostate cancer mortality after 12 years SPCG4 (sse Bill-Axelson point 4.2.1) 12.5% vs 17.9% RD = -5.4% (-11.1 to 0.2); RR = 0.65 (0.45-0.94); p=0.03 Thus, RP is likely to reduce the risks of overall mortality, prostate-cancer mortality and distant metastases compared to WW, but the magnitude of the effect is unclear (width of CI) and the risk reductions appear to have been limited to men less than 65 years of age.</p>	<p>year (SPCG-4): 21.7% vs 45.6%) RD = -23.9% (-30.9 to -16.8); RR = 0.36 (0.27 to 0.47); p< 0.001 Erectile dysfunction after 4 years (SPCG-4): 80% vs 45% RD = 35% (25 to 45) RR = 1.78 (1.48 to 2.15) Urinary leakage after 4 years (SPCG-4): 48.7% vs 21.3% RD = 27% (17 to 37) RR = 2.29 (1.63 to 3.22) Urinary obstruction after 4 years (SPCG-4): 34.5% vs 49.3% RD = -15% (-26 to -4) Bowel symptoms after 4 years (SPCG-4): no ss difference Psychological function after 4 years (SPCG-4): no ss difference</p>	<p>Difference between WW and active surveillance (where patient are monitored closely and where appropriate treatment is promptly initiated) RP appears to increase the risks of erectile dysfunction and urinary leakage but confident statements cannot be made about how frequently these adverse effects occur. Furthermore these estimates must be interpreted cautiously as they are derived from data obtained from a self-administered questionnaire survey of a sample of the trial participants (N = 326), within no baseline quality of life data were obtained and nerve-sparing surgery was not routinely performed on trial participants undergoing RP. No pooling was done</p>
<p>Wilt T.J. & al. "Systematic Review: Comparative Effectiveness and Harms of Treatments for Clinically Localized Prostate Cancer"</p>	<p>Systematic review to determine: - the comparative short- & long-term benefits and harms of therapies for clinically localized prostate cancer - how patients and tumour characteristics affect the outcomes of these therapies. Funding : AHRQ Search date: Mi-Sept 2007 Database: - For RCT : Cochrane</p>	<p>Men with clinically localized prostate cancer (T1 or T2)</p>	<p>Any prostate cancer treatment vs Any prostate cancer treatment</p>	<p>For RP vs WW: Overall survival (VACURG) = 10.6 years for RP vs 8 years for WW (p>0.05) (but underpowered study). All cause mortality after 10 years (SPCG-4) : RD = 5% (-2.8 to 13.0) Prostate cancer mortality after 10 years (SPCG-4): 10% vs 15% (p=0.01)</p>	<p>Distant metastases after 10 years (SPCG-4) : 15.2% vs 25.4%; RD = 10.2% (3.1 to 17.2) Sexual dysfunction (SPCG-4): RR = 1.2 to 18.0 for specific domains Overall distress from all urinary symptoms (SPCG-4): 27% vs 18% ;</p>	<p>High Effectiveness of RP vs WW for overall and disease-specific survival but may be limited to men < 65 y. Increase risk of sexual and urinary dysfunction with RP vs WW ; worse bowel symptom after WW vs RP. Assessment of the comparative effectiveness and harms of localized prostate cancer treatments difficult because of limitations in the evidence.</p>



2008	<p>library, Cochrane Review Group in Prostate Diseases and Urologic malignancies specialized registry (Nov 2007);</p> <ul style="list-style-type: none">- For observational studies (to demonstrate the range of specific outcomes) : PubMed (1991-2004 without update).- For long-term HRQoL: Prostate Cancer Outcomes Study (PCOS)- For outcomes with emerging therapy : Medline (April 2004- Sept 2007) & contact with Endocare- For Effect of patient and tumour characteristics: reviewing RCT, AUA database and U.S. population-based observational studies <p>Included studies:</p> <p>For RP vs. WW: 2 RCT</p> <ol style="list-style-type: none">1. SPCG-4 : Bill-Axelson et al., 2005 (sse Bill-Axelson point 4.2.1) (good quality)2. VACURG: (poor quality) (see Iversen, point 4.2.1)	<p>RR = 1.5 (1.0 to 2.3)</p> <p>Distress from all bowel symptoms (SPCG-4): 3% vs 6%</p> <p>Inability to attain an erection (PCOS): 86% if AD, 58% if RP vs 33% in WW.</p> <p>Patient satisfaction (PCOS): higher with early intervention vs WW.</p>	<p>None RCT enrolled men with primarily PSA-detected disease</p>
-------------	--	---	--



Appendix 5.2. Primary Studies For Treatment

Appendix 5.2.1. RCT : Mortality-morbidity

I Study ID	II Method	III Patient characteristics	IV Intervention(s)	V Results primary outcome	VI Results secondary and other outcome(s)	VII Critical appraisal of study quality
<p>Bill-Axelson et al. "Radical prostatectomy versus Watchful waiting in early prostate cancer"</p> <p>2011</p> <p>(+ Homberg 2002 for design)</p>	<ul style="list-style-type: none"> RCT SPCG-4 Funding: Swedish Cancer Society & National Institutes of Health Setting: 14 centers in Sweden, Finland & Iceland Sample size: 695 Recruitment duration: Oct 1989- Feb 1999 Follow-up: 15 years 	<ul style="list-style-type: none"> Newly diagnosed, localized prostate cancer T0d, T1, T2 (1978 criteria). Tumour well or moderately well differentiated Predominantly detected by symptoms, rather than PSA. Life expectancy >10 years < 75 years old. No other known cancers Serum PSA level < 50 ng/ml. < 25% of the tumour = Gleason 4 & < 5%= Gleason 5. No sign of obstruction of the upper urinary tract. Negative bone scan <p>Comparable groups with the exception of a somewhat higher proportion of men with stage T1b tumours in the WW group; but most of the men had stage T2 tumours.</p>	<p>RP (n=347)</p> <p>= pelvic lymph node evacuation and Walsh-Lepor radical prostatectomy if no nodal metastases found</p> <ul style="list-style-type: none"> hormonal therapy if local recurrence (orchidectomy or gonadotropin-releasing hormones analogues) hormonal therapy if metastases detected by bone scan or, after 2003, if signs of tumour progression, including elevations of PSA level. <p>WW (n=348)</p> <p>= no initial treatment (except TURP already done) :</p> <ul style="list-style-type: none"> transurethral resection if signs of obstructive voiding disorders hormonal therapy if metastases detected by bone scan or, after 2003, if signs of tumour progression, including elevations of PSA level. <p>Follow-up every 6 month (clinical examination and blood test thereafter) for the</p>	<ul style="list-style-type: none"> All cause death : Absolute risk reduction (ARR)=6.6 (-1.3 to 14.5) RR=0.75 (0.61 to 0.92) P=0.007 NNT=15 but for men < 65 years: ARR= 13.5 (2.4 to 24.7) RR=0.52 (0.37 to 0.73) P< 0.001 NNT=7 Prostate cancer death : ARR=6.1 (0.2-12.0) RR=0.62 (0.44-0.87) P=0.01 	<ul style="list-style-type: none"> Distant metastases: ARR=11.7 (4.8 to 18.6) RR=0.59 (0.45 to 0.79) P < 0.001 Local progression : ARR=27.9 (20.9 to 34.8) RR=0.34 (0.26 to 0.45) Post-operative complications after RP: Impotence= 58.1% (52.7 to 64.1) Urinary leakage=32.2%(27.2 to 38.1) 	<ul style="list-style-type: none"> No patients lost to follow-up until December 31, 2009 Critical appraisal: High But Trial initiated before the era of screening by PSA. Radical nature of surgery rather preservation of potency. Population < 75 years with >10 LE (not representative of general population) WW = not Active surveillance (where patients are monitored closely and promptly treated if signs of progression)



			first 2 years, then annually (+bone scan and chest x-ray annually before 2003, bone scan every 2 years after)			
<p>Iversen et al. “Radical prostatectomy versus Expectant treatment for early carcinoma of the prostate” 1995</p>	<ul style="list-style-type: none"> • RCT VACURG • Funding: ? • Setting : 15 Veterans Administration Hospitals in U.S. • Sample size: 111/142 • Recruitment duration: May 1967-March 1975 • Follow-up: 15 y 	<ul style="list-style-type: none"> • Newly diagnosed, histologically confirmed adenocarcinoma of the prostate • Stage I and II (Gleason) • Previously untreated <p>Comparable group a part for mean age (62.7 years in RP vs 66.0 years in EP)</p>	<p>RP</p> <p>= retropubic radical prostatectomy + oral placebo</p> <p>Without prior pelvic lymphadenectomy</p> <p>Expectant Treatment (ET)</p> <p>= oral placebo alone</p> <p>Follow-up every 6 month (no bone scan!) until 1978, no regular FU after because of a lack of funding.</p>	<ul style="list-style-type: none"> • Overall survival: 10.6 vs 8 years (NSS) 	<ul style="list-style-type: none"> • No patients lost to follow-up • Critical appraisal: Low • Trial initiated before the era of screening by PSA • Possible enrollment of patients with disseminated disease because of lack of bone scan and diagnostic lymphadenectomy in routine evaluation. • Small sample size • Drop out (31) 	
<p>Wilt et al . “Radical Prostatectomy versus Observation for Localized Prostate Cancer” 2012</p>	<ul style="list-style-type: none"> • RCT : PIVOT • Funding : Dpt of Veterans Affairs, National Cancer institute and AHRQ • Setting : 44 Department of Veterans Affairs sites and 8 National Cancer Institute sites. • Sample size : 731 (364+367) • Recruitment duration : Nov 1994-Jan2002 	<ul style="list-style-type: none"> • Histologically confirmed, clinically localized prostate cancer (stage T1-T2NxM0 of any grade) diagnosed within the previous 12 months • The mean age was 67 years. 1/3 of the patients • were black; 85% reported full independence in activities of daily living. The median PSA value was • 7.8 ng per milliliter (mean, 10.1). About 50% of the men had stage T1c disease (not palpable, detected by means of PSA testing), and 	<ul style="list-style-type: none"> • RP:technique was let at the surgeon’s discretion • Observation ; Men were offered palliative therapy or chemotherapy for symptomatic or metastatic progression. • Follow-up : every 6 months for a minimum of 8 years and a maximum of 15 years or until the patient died. Bone scans were obtained at 5, 10, and 15 years or at the last visit for persons with less than 15 years of follow-up with additional scans obtained at the 	<ul style="list-style-type: none"> • All-cause mortality : • Among the entire group (HR:; 0.88; 95% confidence interval [CI], 0.71 to 1.08; P = 0.22; absolute risk reduction, 2.9 percentage points; 95% CI, -4.1 to 10.3 in favour of RP). • The absolute reduction in mortality with radical prostatectomy was not significant at any interval and declined over time, from 4.6 	<p>Prostate cancer mortality :</p> <ul style="list-style-type: none"> • Among the entire group (hazard ratio, 0.63; 95% CI, 0.36 to 1.09; P = 0.09; absolute risk reduction, 2.6 percentage points; 95% CI, -1.1 to 6.5) • Among men with PSA< 10 ng/ml : (hazard ratio, 1.03; 95% CI, 0.79 to 1.35). • Among men with PSA > 10 ng/ml : (hazard ratio, 0.67; 95% CI,0.48 to 0.94) • Among men with low risk tumour : (hazard ratio, 1.15; 95% CI, 0.80 to 1.66). • Among men with 	<ul style="list-style-type: none"> • No patients lost to follow-up • Intention to treat analysis • About 50% of the men had disease (not palpable, detected by means of PSA testing. • Critical appraisal : high



<ul style="list-style-type: none"> • Median follow-up : 10 years 	<p>about 25% had histologic scores of 7 or higher on the Gleason scale; 40% of the men had low-risk, 34% intermediate-risk, and 21% high-risk prostate cancer</p> <ul style="list-style-type: none"> • Randomization process was designed to create comparable treatment groups 	<p>clinician's discretion.</p>	<p>percentage points (95% CI, -0.2 to 9.3) at 4 years to 2.9 percentage points (95% CI, -4.2 to 10.0) at 12 years.</p>	<p>intermediate risk tumour : (hazard ratio, 0.69; 95% CI, 0.49 to 0.98; absolute risk reduction, 12.6 percentage points)</p> <ul style="list-style-type: none"> • Among men with high risk tumour:: nonsignificant absolute reduction in mortality of 6.7 percentage points, as compared with observation (P = 0.16)
---	--	--------------------------------	--	--

Appendix 5.2.2. Quality of life

I Study ID	II Method	III Patient characteristics	IV Intervention(s)	VI Results secondary and other outcome(s)	VII Critical appraisal of study quality
<p>Fransson P, et al. Health-related quality of life 10 years after external beam radiotherapy or watchful waiting in patients with localized prostate cancer 2009</p>	<ul style="list-style-type: none"> • RCT • Funding: Swedish Cancer Society (Unmea Trial 1 Study) • Setting:? • Sample size: 54/72 • Recruitment during 1986-1996 • Begin of this follow-up in 2004 • Follow-up: 4 and 10 years <p>According to the authors, the UMEA1 trial is not yet ready for publication!</p> <p>2 questionnaires:</p> <ul style="list-style-type: none"> • PCSS (updated version of the validated QUFW94 formula) • EORTC QLQ-C30 	<ul style="list-style-type: none"> • Localized prostate cancer. • No previous treatment • No other disease with an expected survival time < normal population of the same age <p>Group comparability: ? (appears OK for age but for PSA?)</p>	<p>RT (n=27)</p> <p>5 fractions/week with daily dose of 2.0 Gy.(change in protocol in 1993 from four-field small-box technique to four-field conformal radiation therapy)</p> <p>WW (n=27)</p> <p>= monitoring regularly and treatment deferred until progression.</p>	<ul style="list-style-type: none"> • HRQoL : no statistically significant differences (symptom scale; limitation in daily life; life situation) and few change over time from 4 to 10 year • Symptom evaluation : <ul style="list-style-type: none"> ○ Urinary bother : no difference a part weak urinary stream (mean 4.8 vs 3.0; p=0.034) ○ Bowel symptoms : no difference ○ Sexual bother : more after RT (mean : 7.4 vs 3.8, p=0.011) but no difference in erectile function, nor in maintaining a sufficient erection to perform intercourse, nor in the HRQoL question) 	<ul style="list-style-type: none"> • Decreasing number of questionnaires completed in the FU. • Critical appraisal: Medium <p>Small sample size</p> <p>Lack of baseline data because evaluating HRQoL was not the aim of the study protocol.</p> <p>PCSS instrument validated but maybe difficult to interpret.</p> <p>Quid about PSA detection or not? Quid about the kind of follow-up.</p>



<p>Johansson E, Steineck G et al. Long-term quality-of-life outcomes after radical prostatectomy or watchful waiting.</p>	<ul style="list-style-type: none"> • Longitudinal analysis from a RCT • Funding: Swedish Cancer Society & National Institutes of Health • Setting: 14 centers in Sweden, Finland & Iceland • Sample size: 400 living from SPCG-4; 349 answered once; 166 answered the questionnaire twice. • Recruitment duration: Oct 1989- Feb 1999 • Questionnaires in 1997-1998 and between Oct 2006-Nov 2008 • Follow-up varied from 7 to 17 years (median 12.2) 	<ul style="list-style-type: none"> • 400 living Swedish and Finnish men from the 695 SPCG-4 population (Newly diagnosed, localized prostate cancer; Life expectancy >10 years; < 75 years old (cfr Bill-Axelsson)) • 300 Swedish men from the Swedish Total Population Register, matched for region and age = compared group to understand the effect of leaving the prostate in place 	<p>RP (n=182 but 171 with really RP)</p> <ul style="list-style-type: none"> - hormonal therapy if local recurrence (19%) 	<p>WW (n=167)</p> <ul style="list-style-type: none"> - transurethral resection if signs of obstructive voiding disorders (15%) - hormonal therapy if metastases detected by bone scan or, after 2003, if signs of tumour progression, including elevations of PSA level (28%). 	<p>No intervention because no prostate cancer = controlled group (CG) (n=214)</p>	<ul style="list-style-type: none"> • Number of physical symptoms: No difference RP-WW : 94% vs 94% reported 1 to 4 of erectile dysfunction, weak urinary stream, urinary leakage, or nocturia. But 65% in the CG • Erectile dysfunction (not exhaustive) <ul style="list-style-type: none"> ○ 84% in RP, 80% in WW & 46% in CG. ○ More men in distress due to erection in RP (48%) vs WW (36%) or CG (37%): RR*=1.30 (1.00-1.70) RP vs WW ○ More men in distress from lower self-esteem due to diminished erection in RP (39%) vs WW (23%) or CG (19%): RR*=1.67 (1.20-2.33) RP vs WW ○ Less frequency of orgasm (>1/last 6 months) in RP (18%) vs WW (26%) or CG (58%): RR*=0.62 (0.42-0.91) RP vs WW • Urinary functions (not exhaustive) <ul style="list-style-type: none"> ○ Less weak stream in RP (29%) vs WW (40%) : RR*= 0.71 (0.53-0.96) ○ Less nocturia in RP (49%) vs WW (63%): RR*= 0.79 (0.65-0.95) ○ More urinary leakage at least once daily in RP (41%) vs WW (11%) or CG (3%): RR*= 3.79 (2.36-6.06) RP vs WW ○ More night -time urinary leakage at least once a week in RP (20%) vs WW (8%) or CG (1%): RR*= 2.58 (1.42-4.69) RP vs WW ○ More feeling distress from urinary leakage in RP (18%) vs WW (9%) or CG (4%): RR*= 2.08 (1.15-3.78) RP vs WW ○ More regular dependence on some form of protective aid in RP (54%) vs WW (25%) or CG (8%): RR*=2.15 (1.60-2.90) RP vs WW • QoL <ul style="list-style-type: none"> ○ Lower scores in all psychological measures if RP or WW vs CG but without significant result a part for anxiety. ○ Moderate to high level of anxiety reported by the same proportion of patients in RP (43%) and WW (43%) but by fewer in the CG (33%): RR*=1.42 (1.07-1.88) 	<ul style="list-style-type: none"> • Living and answering portion of the initial sample of SPCG-4. • Critical appraisal: Medium <p>Diagnosis based on clinical symptom (not PSA detected). Time to manifestation of erectile dysfunction or urinary disorder will be substantially longer for men diagnosed as a result of PSA screening.</p> <p>RP more aggressive in the 1990s.</p> <p>About ¼ of men from RP & WW group were AD; thus, perhaps with lower libido (and less acceptance of problem cfr Johansson 2008).</p> <p>Absence of baseline data</p> <p>Questionnaire validated in an unpublished pilot study and in other studies (?)</p>
<p>2011</p>	<p>Face-validated study-specific questionnaire</p> <p>Questionnaire 1997-1998 : 77 items</p> <p>Questionnaire 2006-2008: 141 items</p> <p>Identical question about QoL and functional outcome in both questionnaires.</p> <p>Additional information collected about potential confounders and effect-modifying factors (i.e concurrent diseases and treatments)</p>	<p>Comparability of RP and WW but need of age adjusted calculation to offset the younger age of the compared group (because of a probable error in the age-interval machine).</p>					



*RR**= age adjusted risk relatif

Appendix 5.3. Publications Not Included

Bill-Axelsson A, et al. 2008 Radical prostatectomy versus watchful waiting in localized prostate cancer: the Scandinavian prostate cancer group-4 randomized trial	In Hegarty 2010 SPCG4
Bill-Axelsson A, et al. 2005 Radical prostatectomy versus watchful waiting in early prostate cancer	In Hegarty 2010 SPCG4
Dall'Era MA, et al. 2008 Active surveillance for low-risk prostate cancer: Selection of patients and predictors of progression	Review with a QA = 1 (only Medline, no information about QA of article, data extraction...) Out after QA
Eggerer SE, et al. 2007 Focal therapy for localized prostate cancer: a critical appraisal of rationale and modalities. Review 50 refs	Nothing about watchful waiting or active surveillance Out after text revision (subject)
Frattaroli J, 2008. Clinical events in prostate cancer lifestyle trial: results from two years of follow-up	Out after text revision (subject)
Holmberg L, et al. 2002 A randomized trial comparing radical prostatectomy with watchful waiting in early prostate cancer	In Hegarty 2010 SPCG4
Johansson E, et al. 2009 Time, symptom burden, androgen deprivation, and self-assessed quality of life after radical prostatectomy or watchful waiting:	In Hegarty 2010 SPCG4
Kasperzyk JL, et al. 2011 Watchful waiting and quality of life among prostate cancer survivors in the physicians' health study	Design: Cohort of physicians followed in a clinical trial for CV disease and cancer prevention!
Khatami A, et al. 2006 "PSA doubling time predicts the outcome after active surveillance in screening-detected prostate cancer: results from the European randomized study of screening for prostate cancer, Sweden section." 2007	In Hegarty 2010, Where it is not included because: "This RCT does not randomise men to Radical Prostatectomy and an observation approach" Out : scope (FU of group under surveillance to measure PSA doubling time.
Steineck G, et al. 2002 New England Journal of Medicine 347(11):790-6 - Quality of life after radical prostatectomy or watchful waiting	In Hegarty 2010 SPCG4
Studer UE, et al. 2008 Using PSA to guide timing of androgen deprivation in patients with T0-4 N0-2 M0 prostate cancer not suitable for local curative treatment (EORTC 30891)	Scope : PSA to guide timing of Androgen Deprivation; nor watchfull waiting or active surveillance vs AD not suitable for local curative treatment
Studer UE, et al. 2006 Journal of Clinical Oncology 24(12):1868-76 - Immediate or deferred androgen deprivation for patients with prostate cancer not suitable for local treatment with curative intent: European Organisation for Research and Treatment of Cancer (EORTC) Trial 30891	Idem Studer 2008
Thong MS, et al. 2009 Prostate cancer survivors who would be eligible for active surveillance but were either treated with radiotherapy or managed expectantly: comparisons on long-term quality of life and symptom burden	Design : cross-sectional (level of evidence 2c); 71 men, 10 years after low-risk localized prostate cancer managed with AS (systematic monitoring of men for whom curative treatment is deferred at diagnosis and who receive subsequent curative treatment when the tumour shows progression or when patients decide to change the treatment) were matched with 71 men managed by RT.



	Result : comparable HRQoL and lower symptom burden (bowel function, getting and maintaining an erection) for AS vs RT, even after controlling for comorbidity and disease progress. Out : desing : no RCT
van den Bergh RC, et al. 2010 Is delayed radical prostatectomy in men with low-risk screen-detected prostate cancer associated with a higher risk of unfavorable outcomes?	Included in Hegarty 2010 Where it is not included because it is not a RCT Design: evaluation of a prospective, single arm; observational protocol-based Active Surveillance program Outcome : only biological tumour progression : gleason score, capsular penetration, tumour volume Out : desing : no RCT
ProtecT trial	First results should be published after 2015 (http://www.epi.bris.ac.uk/protect/news/news.htm).
Start Trial	Unfortunately, this study was stopped early due to poor recruitment.No interim reports was published.

Appendix 5.4. Guidelines

Appendix 5.4.1. Patient point of view

Reference	Search date	Recommendations/conclusion	Level of evidence
AUA 2007	2004	Standard: A patient with clinically localized prostate cancer should be informed about the commonly accepted initial interventions including, at a minimum, active surveillance, radiotherapy (external beam and interstitial), and radical prostatectomy. A discussion of the estimates for benefits and harms of each intervention should be offered to the patient.	Panel consensus Low
		Standard: Patient preferences and health conditions related to urinary, sexual, and bowel function should be considered in decision making. Particular treatments have the potential to improve, to exacerbate or to have no effect on individual health conditions in these areas, making no one treatment modality preferable for all patients.	Panel consensus Low
VIKC 2007	2005	Ter ondersteuning van de besluitvorming ten aanzien van de therapiekeuze moet de voorlichting aan patiënten met prostaatacarcinoom kwantitatieve gegevens bevatten over de resultaten en consequenties van de verschillende behandelmogelijkheden van de eigen kliniek.	Panel consensus Low
		Naast de behandelende arts speelt ook een gespecialiseerde verpleegkundige een belangrijke rol bij de voorlichting en begeleiding van mannen met prostaatacarcinoom. Het verdient aanbeveling in de status te vermelden wie de voorlichting heeft gegeven, welke behandelingsmogelijkheden zijn voorgesteld, en wat de te verwachten bijwerkingen zijn.	Panel consensus Low
		Bij de behandeling van patiënten met prostaatkanker is een gestructureerd, multidisciplinair overleg gewenst.	Panel consensus Low
NICE 2008	Until 2007	Men with prostate cancer should be offered individualised information tailored to their own needs. This information should be given by a healthcare professional (for example, a consultant or specialist nurse) and may be supported by written and visual media (for example, slide sets or DVDs).	Low
		Healthcare professionals caring for men with prostate cancer should ascertain the extent to which the man wishes to be involved in decision making and ensure that he has sufficient information to do so.	Low
		A validated, up-to-date decision aid is recommended for use in all urological cancer multidisciplinary teams	high quality evidence and GDG



(MDTs). It should be offered to men with localised prostate cancer when making treatment decisions, by healthcare professionals trained in its use¹ . consensus.

Healthcare professionals should adequately inform men with prostate cancer and their partners or carers about the effects of prostate cancer and the treatment options on their sexual function, physical appearance, continence and other aspects of masculinity. Healthcare professionals should support men and their partners or carers in making treatment decisions, taking into account the effects on quality of life as well as survival. qualitative evidence and GDG consensus.

Appendix 5.4.2. Risk Assessment

Reference	Search date	Recommendations/conclusion	Level of evidence
AUA 2007	2004	Standard: An assessment of the patient's life expectancy, overall health status, and tumour characteristics should be undertaken before any treatment decisions can be made.	Panel consensus Low
VIKC 2007	2005	Bij iedere patiënt met verdenking op prostaatacarcinoom wordt de familie anamnese afgenomen. Als er op basis van de familieanamnese aanwijzingen zijn voor erfelijk prostaatacarcinoom dan wordt periodiek onderzoek verricht volgens de adviezen van de Stichting Opsporing Erfelijke Tumoren. De Gleason score (inclusief de samenstellende componenten) wordt gebruikt bij de gradering van het prostaatacarcinoom en bij de beoordeling van prostaataaldbiopten.	Panel consensus Low Panel consensus Low
NICE 2008	Until June 2007	To help men decide whether to have a prostate biopsy, healthcare professionals should discuss with them their prostate specific antigen (PSA) level, digital rectal examination (DRE) findings (including an estimate of prostate size) and comorbidities, together with their risk factors (including increasing age and black African and Caribbean ethnicity) and any history of a previous negative prostate biopsy. The serum PSA level alone should not automatically lead to a prostate biopsy. Men and their partners or carers should be given information, support and adequate time to decide whether or not they wish to undergo prostate biopsy. The information should include an explanation of the risks (including the increased chance of having to live with the diagnosis of clinically insignificant prostate cancer) and benefits of prostate biopsy. Urological cancer multidisciplinary teams (MDTs) should assign a risk category to all newly diagnosed men with localised prostate cancer.	Low Well designed North American observational studies and GDG consensus Low Low (cohort studies)



Appendix 5.5.1. Treatment

Reference	Search date	Recommendations/conclusion	Level of evidence
AUA 2007	2004	Standard: A patient with clinically localized prostate cancer should be informed about the commonly accepted initial interventions including, at a minimum, active surveillance, radiotherapy (external beam and interstitial), and radical prostatectomy. A discussion of the estimates for benefits and harms of each intervention should be offered to the patient.	Panel consensus Low
		Option: Active surveillance, interstitial prostate brachytherapy, external beam radiotherapy, and radical prostatectomy are appropriate monotherapy treatment options for the patient with low-risk localized prostate cancer.	Panel consensus Low
		Standard: Patient preferences and health conditions related to urinary, sexual, and bowel function should be considered in decision making. Particular treatments have the potential to improve, to exacerbate or to have no effect on individual health conditions in these areas, making no one treatment modality preferable for all patients.	
		Standard: When counseling patients regarding treatment options, physicians should consider the following: Based on outcomes of one randomized controlled clinical trial, when watchful waiting and radical prostatectomy are compared, radical prostatectomy may be associated with a lower risk of cancer recurrence, cancer-related death, and improved survival. (based on SPCG4)	High
VIKC 2007	2005	Bij patiënten met een laag risico (T1c02a, Gleason < 7, PSA < 10 ng/mL) en een gevorderde leeftijd (>75jaar) verdient actief volgen de voorkeur. Daarbij legt men uit dat de levensverwachting niet wordt bepaald door het prostaatacarcinoom en dat elke behandeling een kans heeft op bijwerkingen. Ook bij patiënten met een matig of hoog risico wordt actief volgen verwoogen indien er naast de leeftijd sprake is van duidelijke comorbiditeit die de levensverwachting negatief beïnvloedt.	Intermediate (2+3)
NICE 2008	Until 2007	Men with low-risk localised prostate cancer who are considered suitable for radical treatment should first be offered active surveillance.	Low
		Active surveillance is particularly suitable for a subgroup of men with low-risk localised prostate cancer who have clinical stage T1c, a Gleason score 3+3, a PSA density < 0.15 ng/ml/ml and who have cancer in less than 50% of their total number of biopsy cores with < 10mm of any core involved.	Low
		Active surveillance should be discussed as an option with men who have intermediate-risk localised prostate cancer.	Low
		Active surveillance is not recommended for men with high-risk localised prostate cancer.	Low
EAU 2011 Indications	Until 2010	Jan Inclusion criteria for active surveillance with the lowest risk of cancer progression are: PSA < 10 ng/ml, biopsy Gleason score < 6, < 2 positive biopsies, < 50% cancer per biopsy, cT1c-2a.	Low
		Stage T1a: well and moderately differentiated tumours. In younger patients with a life expectancy of > 10 years, re-evaluation with PSA, TRUS and biopsies of the prostatic remnant is recommended	Moderate (2A)
		Stage T1b-T2b: well and moderately differentiated tumours. In asymptomatic patients with a life expectancy of < 10 years	Moderate (2A)



EAU 2011 Options	Until 2010	Jan	<p>In presumed localised PCa (Nx-N0, M0): Stage T1b-T2b patients who are well informed and have well-differentiated (or Gleason 2-4) PCa and a life expectancy of 10-15 years. All patients not willing to accept side-effects of active treatment. Well-informed, asymptomatic patients with high PSA levels for whom cure is unlikely</p>	Low
Aragon 2008	Until 2007	Nov	<p>In patients with clinically localised prostate cancer with a life expectancy exceeding 10 years, radical prostatectomy or external beam radiotherapy is recommended</p> <hr/> <p>In patients with clinically localised prostate cancer with a life expectancy below 10 years, watchful waiting may be an alternative.</p> <hr/> <p>In patients with clinically localised prostate cancer at low risk, Gleason < 3 + 3, < 50% affected cylinders in the biopsy and PSA < 15 ng/ml, active surveillance can be offered as an alternative to immediate radical treatment.</p>	Moderate (B)
				Moderate (B)
				Very Low (C)



APPENDIX 6. GRADE

Appendix 6.1. Levels of evidence

Quality level	Definition	Methodological Quality of Supporting Evidence
High (A)	We are very confident that the true effect lies close to that of the estimate of the effect	RCTs without important limitations or overwhelming evidence from observational studies
Moderate (B)	We are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different	RCTs with important limitations (inconsistent results, methodological flaws, indirect, or imprecise) or exceptionally strong evidence from observational studies
Low (C)	Our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect	RCTs with very important limitations or observational studies or case series
Very low (D)	We have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of the effect	

Appendix 6.2. Down- or upgrading the evidence

Study Design	Quality of Evidence	Lower if	Higher if
Randomized trial →	High	Risk of bias -1 Serious -2 Very serious	Large effect +1 Large +2 Very large
	Moderate	Inconsistency -1 Serious -2 Very serious	Dose response +1 Evidence of a gradient
Observational study →	Low	Indirectness -1 Serious -2 Very serious	All plausible confounding +1 Would reduce a demonstrated effect or
	Very low	Imprecision -1 Serious -2 Very serious Publication bias -1 Likely -2 Very likely	+1 Would suggest a spurious effect when results show no effect

Appendix 6.3. Strength of recommendations

Appendix 6.3.1. Definitions

Grade	Definition
Strong	The desirable effects of an intervention clearly outweigh the undesirable effects, or clearly do not
Weak	The desirable effects of an intervention probably outweigh the undesirable effects, or probably do not



Appendix 6.4.1. Factors that influence the strength of a recommendation

Factor	Comment
Balance between desirable and undesirable effects	The larger the difference between the desirable and undesirable effects, the higher the likelihood that a strong recommendation is warranted. The narrower the gradient, the higher the likelihood that a weak recommendation is warranted
Quality of evidence	The higher the quality of evidence, the higher the likelihood that a strong recommendation is warranted
Values and preferences	The more values and preferences vary, or the greater the uncertainty in values and preferences, the higher the likelihood that a weak recommendation is warranted
Costs (resource allocation)	The higher the costs of an intervention—that is, the greater the resources consumed—the lower the likelihood that a strong recommendation is warranted

Appendix 6.5. Grade Profile

Appendix 6.5.1. 3.8.4. Men with life expectancy < 10 years

- In patients with localised prostate cancer and life expectancy < 10 years or with important comorbidities watchful waiting with palliative intent is recommended.**

Results	No. of studies	1	2	3	4	5	Reasons for downgrading	GRADE
After a median follow-up of 10 years, RP did not significantly reduce all-cause mortality: HR=0.88 (0.71 to 1.08); P = 0.22 ²⁴	1 (PIVO T)					1	Imprecision : Insufficient sample size	Moderate

1 : limitations of design, 2 :inconsistency, 3 : indirectness, 4 : imprecision, 5 :publication bias

- Strong recommendation based on side effects of radical treatment (see point 3.4.2.for morbidity and point 3.4.3 for quality of life).**

Appendix 6.5.2. Low-risk localised prostate cancer

- In patients with low-risk localised prostate cancer, eligible and opting for a strategy with curative intent, active surveillance can be considered as a management option, taking into account patient preferences and health conditions related to urinary, sexual, and bowel function.**



Results	No. of studies	1	2	3	4	5	Reasons for downgrading	for	GR ADE
After a median follow-up of 10 years, among men with low-risk tumours (n=296), radical prostatectomy increased not significantly all-cause mortality: HR=1.15 (0.80 to 1.66)	1 (PIVOT)			1	1		Indirectness: follow-up in PIVOT trial is described as watchful waiting, not as active surveillance Imprecision : Insufficient sample size (wide CI)		Low

1 : limitations of design, 2 :inconsistency, 3 : indirectness, 4 : imprecision, 5 :publication bias

- **Strong recommendation based on side effects of radical treatment (see point 3.4.2. for morbidity and point 3.4.3 for quality of life).**

Appendix 6.5.3. Intermediate-risk localised prostate cancer

- **In patients with intermediate-risk localised prostate cancer and particularly those with important co-morbidities and life expectancy < 10years, eligible and opting for a strategy with curative intent, active surveillance should be discussed as a management option taking into account patient preferences and health conditions related to urinary, sexual, and bowel function.**

Results	No. of studies	1	2	3	4	5	Reasons for downgrading	GR ADE
After a median follow-up of 10 years, among men with intermediate-risk tumours (n=249), radical prostatectomy reduced significantly all-cause mortality: HR=0.69 (0.49 to 0.98)	1 (PIVOT)			1	1		Indirectness: follow-up in PIVOT trial is described as watchful waiting, not as active surveillance Imprecision : Insufficient sample size	Low

1 : limitations of design, 2 :inconsistency, 3 : indirectness, 4 : imprecision, 5 :publication bias

- **Strong recommendation based on side effects of radical treatment, especially for men with life-expectancy < 10 y (see point 3.4.2. for morbidity and point 3.4.3 for quality of life).**

Appendix 6.5.4. High-risk localised prostate cancer

- **In patients with high-risk localised prostate cancer, active surveillance is not recommended.**



Results	No. of studies	1	2	3	4	5	Reasons for downgrading	GRAD E
After a median follow-up of 10 years, among men with high-risk tumours (n=157), radical prostatectomy reduced not significantly all-cause mortality: HR=0.69 (0.16 to 1.00)	1 (PIVOT)			1	1		Indirectness: follow-up in PIVOT trial is described as watchful waiting, not as active surveillance Imprecision : Insufficient sample size	Low

1 : limitations of design, 2 :inconsistency, 3 : indirectness, 4 : imprecision, 5 :publication bias

Weak recommendation because CI is near 1. Recommendation may change in further study with larger sample size.

Confirmatory biopsy

A biopsy is recommended one year after the diagnosis.

Level of evidence is low because based on results of the three observational studies.^{38, 41, 43} They found that the most cases of disease reclassification seem to follow the confirmatory biopsy.

Strong recommendation, low level of evidence.

Other tests

PSA measurements every six months, clinical examination or MRI every year each, can be considered.

Level of evidence is low because based on conclusions of the two observational studies.^{41, 43} Level of recommendation is weak. Due to the

lack of study focused on active surveillance, we found no effect of tests on outcomes.

Weak recommendation, low level of evidence.

Routine biopsy

- **After the biopsy performed at one year, routine biopsy at years 4, 7 and 10 should be considered.**
- **GCP based on guidelines.**

Life expectancy < 10y

- **After the age of 80 or in case of life expectancy < 10 year, or in case of significant comorbidity development, it's recommended to stop performing routine biopsy and to offer watchful waiting with palliative intent.**

As for 3.8.4

Strong recommendation, moderate level of evidence.

Disease reclassification

- **PSA>10ng/mL, or PSADT< 3years or clinical change or suspicious lesions at mpMRI should be confirmed by additional biopsy.Disease reclassification is achieved after demonstration of an increase in stage or in Grade (Gleason score ≥ 7).**

Level of evidence is low because based on results of the three observational studies.^{38, 41, 43}

- **Strong recommendation, low level of evidence.**
- **Switching to a radical treatment should be reconsidered in case of disease reclassification.**

This recommendation refers to 3.8.5., 3.8.6, 3.8.7.

GCP based on guidelines



APPENDIX 7. LIFE EXPECTANCY TABLE IN BELGIUM

2007-2009 BELGIQUE						
Hommes						
Age révolu (x)	Population observée (px)	Décès observés (dx)	Probabilité de décès (Qx)	Survivants (Lx)	Décès de la table (Dx)	Espérance de vie (Ex)
birth	194.915	696	0,003571	#####	3.571	76,89
0	188.936	162	0,000857	996.429	854	76,66
1	187.982	58	0,000309	995.575	307	75,73
2	186.065	32	0,000172	995.268	171	74,75
3	182.533	41	0,000225	995.096	224	73,76
4	179.585	22	0,000123	994.873	122	72,78
5	178.603	17	0,000095	994.751	95	71,79
6	180.234	19	0,000105	994.656	105	70,80
7	181.236	13	0,000072	994.552	71	69,80
8	182.278	17	0,000093	994.480	93	68,81
9	182.887	11	0,000060	994.387	60	67,82
10	184.413	22	0,000119	994.328	119	66,82
11	185.094	22	0,000119	994.209	118	65,83
12	185.572	31	0,000167	994.091	166	64,83
13	187.990	28	0,000149	993.925	148	63,85
14	193.114	40	0,000207	993.777	206	62,86
15	198.382	62	0,000313	993.571	311	61,87
16	201.313	63	0,000313	993.260	311	60,89

17	200.595	120	0,000598	992.950	594	59,91
18	198.569	131	0,000660	992.356	655	58,94
19	195.885	150	0,000766	991.701	759	57,98
20	194.812	166	0,000852	990.941	844	57,02
21	192.166	179	0,000931	990.097	922	56,07
22	191.744	160	0,000834	989.175	825	55,12
23	192.446	173	0,000899	988.349	888	54,17
24	196.582	145	0,000738	987.461	728	53,22
25	201.539	163	0,000809	986.733	798	52,26
26	205.604	185	0,000900	985.935	887	51,30
27	207.971	175	0,000841	985.047	829	50,34
28	208.002	200	0,000962	984.219	946	49,39
29	207.010	186	0,000899	983.272	883	48,43
30	205.482	183	0,000891	982.389	875	47,48
31	203.957	200	0,000981	981.514	962	46,52
32	205.488	190	0,000925	980.551	907	45,56
33	209.828	228	0,001087	979.645	1.064	44,61
34	217.477	246	0,001131	978.580	1.107	43,65
35	224.352	254	0,001132	977.473	1.107	42,70
36	230.089	248	0,001078	976.367	1.052	41,75
37	232.700	319	0,001371	975.314	1.337	40,79
38	233.329	330	0,001414	973.977	1.378	39,85
39	234.076	382	0,001632	972.600	1.587	38,91
40	237.124	387	0,001632	971.012	1.585	37,97
41	242.726	428	0,001763	969.428	1.709	37,03
42	249.105	484	0,001943	967.718	1.880	36,09
43	252.109	542	0,002150	965.838	2.076	35,16



44	251.012	644	0,002566	963.762	2.473	34,24
45	247.325	622	0,002515	961.289	2.418	33,33
46	244.272	698	0,002857	958.872	2.740	32,41
47	243.014	777	0,003197	956.132	3.057	31,50
48	240.565	877	0,003646	953.074	3.475	30,60
49	237.006	985	0,004156	949.600	3.947	29,71
50	231.702	1.086	0,004687	945.653	4.432	28,83
51	226.716	1.213	0,005350	941.221	5.036	27,96
52	222.007	1.260	0,005675	936.185	5.313	27,11
53	217.690	1.307	0,006004	930.872	5.589	26,26
54	214.051	1.402	0,006550	925.283	6.060	25,42
55	208.464	1.502	0,007205	919.223	6.623	24,58
56	204.465	1.630	0,007972	912.600	7.275	23,76
57	200.828	1.758	0,008754	905.324	7.925	22,95
58	200.446	1.835	0,009155	897.399	8.215	22,14
59	199.070	2.014	0,010117	889.184	8.996	21,34
60	197.564	2.185	0,011060	880.188	9.735	20,56
61	184.988	2.251	0,012168	870.453	10.592	19,78
62	173.174	2.217	0,012802	859.861	11.008	19,02
63	158.468	2.194	0,013845	848.853	11.752	18,26
64	148.160	2.308	0,015578	837.101	13.040	17,51
65	133.864	2.207	0,016487	824.061	13.586	16,78
66	126.861	2.236	0,017626	810.475	14.285	16,05
67	128.951	2.419	0,018759	796.189	14.936	15,33
68	135.635	2.810	0,020717	781.254	16.186	14,61
69	136.289	3.071	0,022533	765.068	17.239	13,91

70	131.924	3.186	0,024150	747.829	18.060	13,22
71	126.058	3.356	0,026623	729.769	19.428	12,54
72	121.978	3.543	0,029046	710.340	20.633	11,86
73	119.103	4.027	0,033811	689.708	23.320	11,20
74	117.164	4.333	0,036982	666.388	24.645	10,58
75	114.160	4.654	0,040767	641.743	26.162	9,97
76	110.377	5.144	0,046604	615.581	28.688	9,37
77	102.704	5.318	0,051780	586.893	30.389	8,80
78	94.011	5.375	0,057174	556.503	31.818	8,26
79	85.112	5.487	0,064468	524.686	33.825	7,73
80	77.839	5.438	0,069862	490.860	34.293	7,22
81	71.493	5.645	0,078959	456.568	36.050	6,73
82	64.973	5.771	0,088822	420.518	37.351	6,26
83	58.125	5.847	0,100594	383.167	38.544	5,82
84	50.637	5.635	0,111282	344.623	38.350	5,42
85	43.632	5.409	0,123969	306.272	37.968	5,04
86	36.941	5.171	0,139980	268.304	37.557	4,68
87	28.413	4.231	0,148911	230.747	34.361	4,36
88	19.245	3.240	0,168355	196.386	33.063	4,03
89	12.204	2.231	0,182809	163.324	29.857	3,75
90	8.614	1.680	0,195031	133.467	26.030	3,47
91	7.479	1.696	0,226768	107.436	24.363	3,19
92	6.560	1.574	0,239939	83.073	19.933	2,98
93	5.301	1.412	0,266365	63.141	16.818	2,77
94	3.877	1.100	0,283725	46.322	13.143	2,59
95	2.589	788	0,304365	33.179	10.099	2,42



96	1.690	605	0,357988	23.081	8.263	2,26
97	1.048	330	0,314885	14.818	4.666	2,24
98	669	223	0,333333	10.152	3.384	2,04
99	411	167	0,406326	6.768	2.750	1,80
100	218	90	0,412844	4.018	1.659	1,69
101	120	52	0,433333	2.359	1.022	1,53
102	70	29	0,414286	1.337	554	1,33
103	39	23	0,589744	783	462	0,91
104+	19	9	1,000000	321	321	0,17

Cette table de mortalité a été revue et corrigée pour mieux correspondre au type de quotients utilisés (une note explicative complète sur ce sujet est en préparation).

Les changements principaux concernent :

- la série des âges (qui commence par «birth», représentant l'âge exact 0, se poursuit par les âges révolus et se clôture par l'âge «104+») et
- le calcul de l'espérance de vie (moyenne arithmétique des âges au décès de la table à partir d'un âge donné, diminuée du nombre d'années déjà vécues pour atteindre cet âge).

SOURCE : SPF Économie - Direction générale Statistique et Information économique.

<http://statbel.fgov.be>

APPENDIX 8. QUALITY CRITERIA FOR PROSTATE BIOPSY

We put here some important discussions around quality criteria for prostate biopsy.

HGPIN

Question : it remains unclear if and how many nucleoli are needed to diagnose HGPIN (Egevad et al. Mod Pathol 19:180).

Answer : In my opinion, this point is out of the subject of "Active surveillance in prostate adenocarcinoma".

The diagnostic criteria for prostatic intraepithelial neoplasia (LGPIN and HGPIN) are quite well described by D. Bostwick in the Urologic Surgical Pathology 2nd Edition.

Because of its lack of specificity, it is recommended that LGPIN should not be reported (Strigley JR et al., Arch Pathol Lab Med 2000). As noted by J. Epstein and M. Herawi in Journal of Urology 2006 and by others, only multifocal HGPIN seems to be associated with an increase risk of cancer on repeat biopsy.

Comment : The categories of diseases 3 & 4. High grade prostatic intraepithelial neoplasia (HGPIN) and atypical glands suspicious for cancer (ASAP) should never be diagnosed as cancer and never treated as cancer by the clinician.

Extent of tumour involvement of cores

Question: if tumour is discontinuously present in a core, do we suggest to collapse the tumour by disregarding intervening stroma or not? (see for example Karram et al. Am J Surg Pathol 35:1351).

Answer: Currently, there is no evidence one method is better than the other, but it might be relevant to put forward one method, so the same method will be used throughout Belgium (in my opinion, most non-urologists are even not aware of the existence of two systems).

I agree with the remark. As said by Fine et al European Urology 2012, evidences are too limited to draw a definitive conclusion about the method to use. However, the study recently published by Karram S et al. (Am J Surg Pathol 2011) demonstrated that for prostate cancer in which the



needle biopsy grade is representative of the entire tumour, quantifying cancer extent on biopsy by measuring discontinuous cancer on biopsy from one end to the other as opposed to "collapsing" the cancer by subtracting out the intervening benign prostate tissue correlates better with organ-confined disease and risk of positive margins. For this reason, I added the reference in the text.

Perineural and lymphovascular invasion

Question: there is no international consensus regarding the reporting of these features in needle biopsies; especially for perineural invasion there is no agreement. The ISUP consensus on prostatectomies suggested that lymphovascular invasion should be reported, but there is no guideline for perineural invasion on prostatectomies (see series of articles in January 2010 issue of Mod Pathol). Findings on prostatectomy studies cannot be extrapolated to needle biopsies (see Freeman. Surg Oncol 18:200) and several studies show that the clinical relevance of reporting perineural invasion on needle biopsies is still to be proven (Al-Hussain et al. J Urol 186:470, Harnden et al. Cancer 109:13).

Answer: I agree with the remark. The sentence: "Perineural and lymphovascular invasion are currently not considered as essential reporting elements for prostatic needle biopsies by leading international urological pathologists" is added in the text.

Gleason grading

Comments : this is of course pivotal and nicely addressed by the already included reference (ISUP 2005 modified Gleason). Special attention has to be given regarding the accuracy of reporting Gleason patterns. While undergrading has been an issue, it appears that currently there is a tendency to overgrade (see for example Egevad et al. Histopathology, accepted article). There exist several tools to train Gleason grading, e.g. Web-tutorials (<http://162.129.103.34/prostate/index.htm#consensus>). The website might be included in specific recommendation for pathologists or in reference. The current text refers to primary and secondary patterns; it might be considered to add a statement that the 2005 ISUP guidelines

should also be followed in case of minor secondary patterns and tertiary patterns (especially since the latter is differently reported on needle and prostatectomy specimens). Furthermore, the 2005 guidelines could not reach a conclusion regarding grading of glomeruloid structures; more recent data suggest that they should be graded as Gleason 4 (Lotan et al. Hum Pathol 40:471). Recommendations published by Fine SW et al. (2012) for special Gleason grading scenarios such as the context of abundant high-grade cancer, prostate cancer variants, glomeruloid structures and/or the presence of a tertiary Gleason pattern in prostate biopsies (Fine SW et al. European Urology 2012) is now added in the text.

APPENDIX 9. EXPERTS AND STAKEHOLDERS FINAL REVIEW

Letter to stakeholders in preparation of stakeholders meeting:

Instructions for scoring the recommendations

Please score each recommendation on a 5-point scale to indicate your agreement with the recommendation: a score of '1' indicates 'completely disagree', '2' indicates 'somewhat disagree', '3' indicates 'unsure', '4' indicates 'somewhat agree', and '5' indicates 'completely agree'. You can also answer 'not applicable' (NA) in case you are not familiar with the underlying evidence.

In case you disagree with the recommendation (score '1' or '2'), please provide the scientific evidence (which of course should at least be better than the presented evidence).

The scores should be sent back by Tuesday November, 20th at the latest to francoise.mambourg@kce.fgov.be. Scores received afterwards will not be taken into account. The scores will then be anonymised and summarised into a median score, range and % of 'agree'-scores. The scores will be presented at the open meeting on Tuesday November 27th, 18.00-20.00h at the KCE, and serve to focus the discussion.



Item	Recommendations before stakeholders meeting	Recommendations after stakeholders meeting	GOR	LoE	5	4	% 4 or 5
Chapter 3. WATCHFUL WAITING / ACTIVE SURVEILLANCE							
3.8.1	A guideline with criteria of a good quality biopsy (as well for TRUS-guided as for MRI-guided biopsy) is needed not only for the pathologists but also for the urologists	A guideline with quality criteria of a prostate biopsy is needed not only for pathologists but also for urologists and radiologists	GCP		10	2	100%
3.8.2	An assessment should be undertaken before any treatment decisions can be made. This assessment should include: (1) the patient's overall health status, his life expectancy and comorbidities and (2) the quality of the biopsy and tumour characteristics (including the risk category)	An assessment should be undertaken before any treatment decisions can be made. This assessment should include: (1) the patient's overall health status, his life expectancy and comorbidities and (2) the quality of the biopsy and tumour characteristics (including the risk category)	GCP		7	4	92%
3.8.3	A patient with clinically localised prostate cancer, eligible and opting for a strategy with curative intent, should be informed about commonly accepted initial managements with regards to his health status, life expectancy and tumour risk category. Commonly accepted initial managements include at least active surveillance, radiotherapy (external beam and interstitial), and radical prostatectomy. The estimated benefits and harms of each intervention should be explained and discussed with the patient.	A patient with clinically localised prostate cancer, eligible and opting for a strategy with curative intent, should be informed about commonly accepted initial managements with regards to his health status, life expectancy and tumour risk category. Commonly accepted initial managements include at least active surveillance, radiotherapy (external beam and interstitial), and radical prostatectomy. The estimated benefits and harms of each intervention should be explained and discussed with the patient.	GCP		11	0	100%
3.8.4	In patients with localised prostate cancer and life expectancy < 10 years or with important comorbidities watchful waiting with palliative intent is recommended.	In patients with localised prostate cancer and life expectancy < 10 years or with important comorbidities, watchful waiting with palliative intent is recommended.	Strong	Moderate	2	4	60%
3.8.5 - 1	In patients with low-risk localised prostate cancer, eligible and opting for a strategy with curative intent, active surveillance can be considered as a management option, taking into account patient preferences and health conditions related to urinary, sexual, and bowel function.	In patients with low-risk localised prostate cancer, eligible and opting for a strategy with curative intent, active surveillance should be considered as a management option, taking into account patient preferences and health conditions related to urinary, sexual, and bowel function.	Strong	Low	8	2	100%
3.8.5 - 2	Men with low-risk localised prostate cancer must be informed that at the present time there is no demonstrated benefit to be immediately treated.	Men with low-risk localised prostate cancer must be informed that at the present time there is no demonstrated benefit for immediate treatment.	GCP		5	1	60%



3.8.6	In patients with intermediate-risk localised prostate cancer and particularly those with important co-morbidities and life expectancy <10years, eligible and opting for a strategy with curative intent, active surveillance should be discussed as a management option taking into account patient preferences and health conditions related to urinary, sexual, and bowel function.	Because of the heterogeneity of the patients with intermediate-risk localised prostate cancer, no general recommendation can currently be made on active surveillance in this subset of patients.	Strong	Low	4	3	70%
3.8.7	In patients with high-risk localised prostate cancer, active surveillance is not recommended.	In patients with high-risk localised prostate cancer, active surveillance is not recommended.	Weak	Low	8	2	100%
4.6.1	A repeat biopsy is recommended one year after the diagnosis.	A repeat biopsy is recommended one year after the diagnosis.	Strong	Low	7	4	100%
4.6.2	PSA measurements every six months and clinical examination or MRI every year each, can be considered.	PSA measurements and clinical examination every six months can be considered. Imaging each year can be considered.	Weak	Low	5	6	92%
4.6.3	After the biopsy performed at one year, routine biopsy at years 4, 7 and 10 should be considered	After the biopsy performed at one year, routine biopsy at years 4, 7 and 10 can be considered	GCP		1	6	70%
4.6.4	After the age of 80, or in case of life expectancy <10 year, or in case of the development of significant comorbidity it is recommended to stop performing routine biopsies and to offer watchful waiting with palliative intent.	After the age of 80, or in case of life expectancy <10 year, or in case of the development of significant comorbidity it is recommended to stop performing routine biopsies and to offer watchful waiting with palliative intent.	Strong	Moderate	7	4	92%
4.6.5 -1	Disease progress as suggested by PSA>10ng/mL, or PSADT< 3 years , or clinical change, or suspicious lesions at mpMRI, should be confirmed by an additional biopsy. Disease reclassification is achieved after demonstration of an increase in stage or in Grade (Gleason score ≥7).	Disease progression as suggested by PSA>10ng/mL, or PSADT< 3 years, or clinical change, or suspicious lesions at imaging, should be confirmed by an additional biopsy and followed by risk reclassification.	Strong	Low	6	4	90%
4.6.5 - 2	Switching to a radical treatment should be considered in case of disease reclassification.	Switching to a radical treatment should be considered in case of risk reclassification.	GCP		7	4	100%



■ REFERENCES

1. Iversen P, Madsen PO, Corle DK. Radical prostatectomy versus expectant treatment for early carcinoma of the prostate. Twenty-three year follow-up of a prospective randomized study. *Scand J Urol Nephrol Suppl.* 1995;172:65-72.
2. Heindenreich A. Guidelines on Prostate Cancer. European Association of Urology; 2012.
3. Mambourg F, Van den Bruel A, Devriese S, Leys M, Vinck I, Lona M, et al. Health Technology Assessment prostate-specific-antigen (PSA) voor prostaatkankerscreening. Health Technology Assessment (HTA). Brussel: Federaal Kenniscentrum voor de Gezondheidszorg (KCE); 2006 15/05/2006. KCE reports 31A (D2006/10.273/17) Available from: <http://kce.fgov.be>
4. Fervers B, Burgers JS, Haugh MC, Latreille J, Mlika-Cabanne N, Paquet L, et al. Adaptation of clinical guidelines: literature review and proposition for a framework and procedure. *Int J Qual Health Care.* 2006;18(3):167-76.
5. AGREE. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care.* 2003;12(1):18-23.
6. Hegarty J, Beirne PV, Walsh E, Comber H, Fitzgerald T, Wallace Kazer M. Radical prostatectomy versus watchful waiting for prostate cancer. Review. *Cochrane Database of Systematic Reviews.* 2010;11.
7. Parker C. Active surveillance: towards a new paradigm in the management of early prostate cancer. *Lancet Oncol.* 2004;5(2):101-6.
8. Richtlijn Prostaatkarcinoom: diagnostiek en behandeling. Vereniging van Integrale Kankercentra; 2007.
9. NICE. Prostate cancer: diagnosis and treatment. London: National Collaborating Centre for Cancer; 2008.
10. Thompson I, Thrasher JB, Aus G, Burnett AL, Canby-Hagino ED, Cookson MS, et al. Guideline for the management of clinically localized prostate cancer: 2007 update. *J Urol.* 2007;177(6):2106-31.



11. Working group of the Clinical Practice Guideline on Prostate Cancer Treatment. Clinical Practice Guidelines on Prostate Cancer Treatment. . Madrid: Aragon Institute of Health Sciences (I+CS); 2008.
12. Bill-Axelson A, Holmberg L, Ruutu M, Garmo H, Stark JR, Busch C, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *New England Journal of Medicine*. 2011;364(18):1708-17.
13. Wilt TJ, MacDonald R, Rutks I, Shamlivan TA, Taylor BC, Kane RL. Systematic review: comparative effectiveness and harms of treatments for clinically localized prostate cancer. Review 78 refs Erratum appears in *Ann Intern Med*. 2008 Jun 3;148(11):888. *Annals of Internal Medicine*. 2008;148(6):435-48.
14. Hoffman RM, Hunt WC, Gilliland FD, Stephenson RA, Potosky AL. Patient satisfaction with treatment decisions for clinically localized prostate carcinoma. Results from the Prostate Cancer Outcomes Study. *Cancer*. 2003;97(7):1653-62.
15. Dahabreh IJ, Chung M, Balk EM, Yu WW, Mathew P, Lau J, et al. Active surveillance in men with localized prostate cancer: a systematic review. *Ann Intern Med*. 2012;156(8):582-90.
16. Comité d'évaluation des pratiques médicales en matière de médicaments. Traitements efficaces dans les pathologies bénignes et malignes de la prostate. Bruxelles: 2011. Réunions de consensus - Rapports du jury Available from: <http://www.inami.fgov.be/drug/fr/statistics-scientific-information/consensus/index.htm>
17. Johansson E, Steineck G, Holmberg L, Johansson JE, Nyberg T, Ruutu M, et al. Long-term quality-of-life outcomes after radical prostatectomy or watchful waiting: the Scandinavian Prostate Cancer Group-4 randomised trial. *Lancet Oncol*. 2011;12(9):891-9.
18. Fransson P, Damber JE, Widmark A. Health-related quality of life 10 years after external beam radiotherapy or watchful waiting in patients with localized prostate cancer. *Scandinavian Journal of Urology & Nephrology*. 2009;43(2):119-26.
19. Holmberg L, Bill-Axelson A, Helgesen F, Salo JO, Folmerz P, Haggman M, et al. A randomized trial comparing radical prostatectomy with watchful waiting in early prostate cancer. *New England Journal of Medicine*. 2002;347(11):781-9.
20. Wilt TJ, Brawer MK, Barry MJ, Jones KM, Kwon Y, Gingrich JR, et al. The Prostate cancer Intervention Versus Observation Trial:VA/NCI/AHRQ Cooperative Studies Program #407 (PIVOT): design and baseline results of a randomized controlled trial comparing radical prostatectomy to watchful waiting for men with clinically localized prostate cancer. *Contemporary Clinical Trials*. 2009;30(1):81-7.
21. Avery KN, Blazeby JM, Lane JA, Neal DE, Hamdy FC, Donovan JL. Decision-making about PSA testing and prostate biopsies: a qualitative study embedded in a primary care randomised trial. *Eur Urol*. 2008;53(6):1186-93.
22. Klotz L. Active surveillance with selective delayed intervention: using natural history to guide treatment in good risk prostate cancer. *J Urol*. 2004;172(5 Pt 2):S48-50; discussion S-1.
23. Bill-Axelson A, Holmberg L, Ruutu M, Haggman M, Andersson SO, Bratell S, et al. Radical prostatectomy versus watchful waiting in early prostate cancer. *N Engl J Med*. 2005;352(19):1977-84.
24. Wilt TJ, Brawer MK, Jones KM, Barry MJ, Aronson WJ, Fox S, et al. Radical prostatectomy versus observation for localized prostate cancer. *N Engl J Med*. 2012;367(3):203-13.
25. Fransson P, Lund JA, Damber JE, Klepp O, Wiklund F, Fossa S, et al. Quality of life in patients with locally advanced prostate cancer given endocrine treatment with or without radiotherapy: 4-year follow-up of SPCG-7/SFUO-3, an open-label, randomised, phase III trial. *Lancet Oncol*. 2009;10(4):370-80.
26. Heidenreich A, Bellmunt J, Bolla M, Joniau S, Mason M, Matveev V, et al. EAU guidelines on prostate cancer. Part 1: screening, diagnosis, and treatment of clinically localised disease. *European Urology*. 2011;59(1):61-71.
27. Botswick D, Cheng L. *Urologic Surgical Pathology*. 2nd Edition ed. London: Elsevier; 2008.



28. Karram S, Trock BJ, Netto GJ, Epstein JI. Should intervening benign tissue be included in the measurement of discontinuous foci of cancer on prostate needle biopsy? Correlation with radical prostatectomy findings. *Am J Surg Pathol*. 2011;35(9):1351-5.
29. Epstein JI, Allsbrook WC, Jr., Amin MB, Egevad LL. The 2005 International Society of Urological Pathology (ISUP) Consensus Conference on Gleason Grading of Prostatic Carcinoma. *Am J Surg Pathol*. 2005;29(9):1228-42.
30. Fine SW, Amin MB, Berney DM, Bjartell A, Egevad L, Epstein JI, et al. A contemporary update on pathology reporting for prostate cancer: biopsy and radical prostatectomy specimens. *Eur Urol*. 2012;62(1):20-39.
31. Hodge KK, McNeal JE, Terris MK, Stamey TA. Random systematic versus directed ultrasound guided transrectal core biopsies of the prostate. *J Urol*. 1989;142(1):71-4; discussion 4-5.
32. Dickinson L, Ahmed HU, Allen C, Barentsz JO, Carey B, Futterer JJ, et al. Magnetic resonance imaging for the detection, localisation, and characterisation of prostate cancer: recommendations from a European consensus meeting. *Eur Urol*. 2011;59(4):477-94.
33. Moore CM, Robertson NL, Arsanious N, Middleton T, Villers A, Klotz L, et al. Image-Guided Prostate Biopsy Using Magnetic Resonance Imaging-Derived Targets: A Systematic Review. *Eur Urol*. 2012.
34. Barentsz JO, Richenberg J, Clements R, Choyke P, Verma S, Villeirs G, et al. ESUR prostate MR guidelines 2012. *Eur Radiol*. 2012;22(4):746-57.
35. Portalez D, Mozer P, Cornud F, Renard-Penna R, Misrai V, Thoulouzan M, et al. Validation of the European Society of Urogenital Radiology Scoring System for Prostate Cancer Diagnosis on Multiparametric Magnetic Resonance Imaging in a Cohort of Repeat Biopsy Patients. *Eur Urol*. 2012.
36. Moore CM, Robertson NL, Arsanious N, Middleton T, Villers A, Klotz L, et al. Reply from Authors re: Behfar Ehdai, Shahrokh F. Shariat. Magnetic Resonance Imaging-targeted Prostate Biopsy: Back to the Future. *Eur Urol*. In press. <http://dx.doi.org/10.1016/j.eururo.2012.06.049>: What Will It Cost to Target Clinically Relevant Prostate Cancer? *Eur Urol*. 2012.
37. NICE. Review of Clinical Guideline (CG58) –Prostate cancer: diagnosis and treatment. In. Review proposal consultation document ed. London: NICE; 2011.
38. Bangma CH, Bul M, Roobol M. The Prostate cancer Research International: Active Surveillance study. *Current Opinion in Urology*. 2012;22(3):216-21.
39. Coen JJ, Feldman AS, Smith MR, Zietman AL. Watchful waiting for localized prostate cancer in the PSA era: what have been the triggers for intervention? *BJU International*. 2011;107(10):1582-6.
40. Ischia JJ, Pang CY, Tay YK, Suen CFDLW, Aw HC, Frydenberg M. Active surveillance for prostate cancer: an Australian experience. *BJU International*. 2012;109 Suppl 3:40-3.
41. Klotz L. Active surveillance: the Canadian experience. *Current Opinion in Urology*. 2012;22(3):222-30.
42. Kravchick S, Peled R, Cytron S. Watchful waiting and active surveillance approach in patients with low risk localized prostatic cancer: an experience of out-patients clinic with 12-year follow-up. *Pathology Oncology Research*. 2011;17(4):893-7.
43. Tosoian JJ, Trock BJ, Landis P, Feng Z, Epstein JI, Partin AW, et al. Active surveillance program for prostate cancer: an update of the Johns Hopkins experience. *Journal of Clinical Oncology*. 2011;29(16):2185-90.
44. Loblaw A, Zhang L, Lam A, Nam R, Mamedov A, Vesprini D, et al. Comparing prostate specific antigen triggers for intervention in men with stable prostate cancer on active surveillance. *J Urol*. 2010;184(5):1942-6.
45. Bul M, Zhu X, Rannikko A, Staerman F, Valdagni R, Pickles T, et al. Radical prostatectomy for low-risk prostate cancer following initial active surveillance: results from a prospective observational study. *Eur Urol*. 2012;62(2):195-200.



46. Ross AE, Loeb S, Landis P, Partin AW, Epstein JI, Kettermann A, et al. Prostate-specific antigen kinetics during follow-up are an unreliable trigger for intervention in a prostate cancer surveillance program. *J Clin Oncol.* 2010;28(17):2810-6.
47. Adamy A, Yee DS, Matsushita K, Maschino A, Cronin A, Vickers A, et al. Role of prostate specific antigen and immediate confirmatory biopsy in predicting progression during active surveillance for low risk prostate cancer. *J Urol.* 2011;185(2):477-82.

